




Europäisches
Patentamt

European
Patent Office

Office européen
des brevets



Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03027637.2

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk



Anmeldung Nr:
Application no.: 03027637.2
Demande no:

Anmeldetag:
Date of filing: 02.12.03
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

BASF AKTIENGESELLSCHAFT

67056 Ludwigshafen
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Clp-protease as target for herbicides

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)

Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

C12Q1/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT RO SE SI SK TR LI

Clp-protease as target for herbicides

5 The present invention relates to Clp-protease, which, when absent, brings about reduced growth and chlorotic leaves as target for herbicides. For this purpose, novel nucleic acid sequences encompassing SEQ ID NO:3, SEQ ID NO:11 and SEQ ID NO: 17 and functional equivalents of SEQ ID NO:3, SEQ ID NO:11 and SEQ ID NO: 17 are provided. Moreover, the present invention relates to the use of Clp-protease in a method for identifying compounds with herbicidal or growth-regulatory activity, and to the use of the compounds identified by this method as herbicides or growth regulators.

10 The basic principle of identifying herbicides via the inhibition of a defined target is known (for example US 5,187,071, WO 98/33925, WO 00/77185). In general, there is a great demand for the detection of enzymes which might constitute novel targets for herbicides. The reasons are resistance problems which occur with herbicidal active ingredients which act on known targets, and the ongoing endeavor to identify novel herbicidal active ingredients which are distinguished by as wide as possible a spectrum of action, ecological and toxicological acceptability and/or low application rates.

20 In practice, the detection of novel targets entails great difficulties since the inhibition of an enzyme which forms part of a metabolic pathway frequently has no further effect on the growth of the plant. This may be attributed to the fact that the plant switches to alternative metabolic pathways whose existence is not known or that the inhibited enzyme is not limiting for the metabolic pathway. Furthermore, plant genomes are distinguished by a high degree of functional redundancy. Functionally equivalent enzymes are found more frequently in gene families in the Arabidopsis thaliana genome than in insects or mammals (Nature, 2000, 408(6814):796-815). This hypothesis is confirmed experimentally by the fact that comprehensive gene knock-out programs by T-DNA or transposon insertion into Arabidopsis yielded fewer manifested phenotypes to date than expected (Curr. Op. Plant Biol. 4, 2001, pp.111-117).

30 It is an object of the present invention to identify novel targets which are essential for the growth of plants or whose inhibition leads to reduced plant growth, and to provide methods which are suitable for identifying herbicidally active and/or growth-regulatory compounds.

35 We have found that this object is achieved by the use of nuclear encoded Clp-protease in a method for identifying herbicides.

40 Further terms used in the description are now defined at this point.

"Affinity tag": this refers to a peptide or polypeptide whose coding nucleic acid sequence can be fused to the nucleic acid sequence according to the invention either

2

directly or by means of a linker, using customary cloning techniques. The affinity tag serves for the isolation, concentration and/or selective purification of the recombinant target protein by means of affinity chromatography from total cell extracts. The above-mentioned linker can advantageously contain a protease cleavage site (for example for thrombin or factor Xa), whereby the affinity tag can be cleaved from the target protein when required. Examples of common affinity tags are the "His tag", for example from Qiagen, Hilden, "Strep tag", the "Myc tag" (Invitrogen, Carlsberg), the tag from New England Biolabs which consists of a chitin-binding domain and an intein, the maltose-binding protein (pMal) from New England Biolabs, and what is known as the CBD tag from Novagen. In this context, the affinity tag can be attached to the 5' or the 3' end of the coding nucleic acid sequence with the sequence encoding the target protein.

"Activity of nuclear encoded Clp-protease": the term activity describes the ability of an enzyme to convert a substrate into a product. The enzymatic activity can be determined in what is known as an activity assay via the increase in the product, the decrease in the substrate (or starting material) or the decrease in a specific cofactor, or via a combination of at least two of the abovementioned parameters, as a function of a defined period of time. "Activity of nuclear encoded Clp-protease" describes here the ability of an enzyme to catalyze the hydrolysis of peptides of maximal five amino acids in vitro.

"Expression cassette": an expression cassette contains a nucleic acid sequence according to the invention linked operably to at least one genetic control element, such as a promoter, and, advantageously, a further control element, such as a terminator. The nucleic acid sequence of the expression cassette can be for example a genomic or complementary DNA sequence or an RNA sequence, and their semisynthetic or fully synthetic analogs. These sequences can exist in linear or circular form, extrachromosomally or integrated into the genome. The nucleic acid sequences in question can be synthesized or obtained naturally or contain a mixture of synthetic and natural DNA components, or else consist of various heterologous gene segments of various organisms.

Artificial nucleic acid sequences are also suitable in this context as long as they make possible the expression, in a cell or an organism, of a polypeptide with the enzymatic activity of a nuclear encoded Clp Protease, preferably with the biological activity of a nuclear encoded Clp Protease, which polypeptide is encoded by a nucleic acid sequence according to the invention. For example, synthetic nucleotide sequences can be generated which have been optimized with regard to the codon usage of the organisms to be transformed.

All of the abovementioned nucleotide sequences can be generated from the nucleotide units by chemical synthesis in the manner known per se, for example by fragment con-

- densation of individual overlapping complementary nucleotide units of the double helix. Oligonucleotides can be synthesized chemically for example in the manner known per se using the phosphoamidite method (Voet, Voet, 2nd Edition, Wiley Press New York, pp. 896-897). When preparing an expression cassette, various DNA fragments can be
- 5 manipulated in such a way that a nucleotide sequence with the correct direction of reading and the correct reading frame is obtained. The nucleic acid fragments are linked with each other via general cloning techniques as are described, for example, in T. Maniatis, E.F. Fritsch and J. Sambrook, "Molecular Cloning: A Laboratory Manual", Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1989), in T.J. Silhavy, M.L.
- 10 Berman and L.W. Enquist, Experiments with Gene Fusions, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1984) and in Ausubel, F.M. et al., "Current Protocols in Molecular Biology", Greene Publishing Assoc. and Wiley-Interscience (1994).
- 15 "Operable linkage" or "functional linkage": an operable, or functional, linkage is understood as meaning the sequential arrangement of regulatory sequences or genetic control elements in such a way that each of the regulatory sequences, or each of the genetic control elements, can fulfill its intended function when the coding sequence is expressed.
- 20 "Functional equivalents" describe, in the present context, nucleic acid sequences which hybridize under standard conditions with the nucleic acid sequence SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15 or SEQ ID NO:17 or parts of the aforementioned nucleic acid sequences and which are capable of bringing about the expression, in a cell
- 25 or an organism, of a polypeptide with the activity of Clp protease.
- To carry out the hybridization, it is advantageous to use short oligonucleotides with a length of approximately 10-50 bp, preferably 15-40 bp, for example of the conserved or other regions, which can be determined in the manner with which the skilled worker is
- 30 familiar by comparisons with other related genes. However, longer fragments of the nucleic acids according to the invention with a length of 100-500 bp, or the complete sequences, may also be used for hybridization. Depending on the nucleic acid/oligonucleotide used, the length of the fragment or the complete sequence, or depending on which type of nucleic acid, i.e. DNA or RNA, is being used for the hybridiza-
- 35 tion, these standard conditions vary. Thus, for example, the melting temperatures for DNA:DNA hybrids are approximately 10°C lower than those of DNA:RNA hybrids of the same length.
- Standard hybridization conditions are to be understood as meaning, depending on the
- 40 nucleic acid, for example temperatures of between 42 and 58°C in an aqueous buffer solution with a concentration of between 0.1 and 5 x SSC (1 x SSC = 0.15 M NaCl, 15 mM sodium citrate, pH 7.2) or additionally in the presence of 50% formamide, such as,

- for example, 42°C in 5 x SSC, 50% formamide. The hybridization conditions for DNA:DNA hybrids are advantageously 0.1 x SSC and temperatures of between approximately 20°C and 65°C, preferably between approximately 30°C and 45°C. In the case of DNA:RNA hybrids, the hybridization conditions are advantageously 0.1 x SSC and temperatures of between approximately 30°C and 65°C, preferably between approximately 45°C and 55°C. These hybridization temperatures which have been stated are melting temperature values which have been calculated by way of example for a nucleic acid with a length of approx. 100 nucleotides and a G + C content of 50% in the absence of formamide. The experimental conditions for DNA hybridization are described in relevant textbooks of genetics such as, for example, in Sambrook et al., "Molecular Cloning", Cold Spring Harbor Laboratory, 1989, and can be calculated using formulae with which the skilled worker is familiar, for example as a function of the length of nucleic acids, the type of the hybrids or the G + C content. The skilled worker will find further information on hybridization in the following textbooks: Ausubel et al. (eds), 1985, "Current Protocols in Molecular Biology", John Wiley & Sons, New York; Hames and Higgins (eds.), 1985, "Nucleic Acids Hybridization: A Practical Approach", IRL Press at Oxford University Press, Oxford; Brown (ed.), 1991, "Essential Molecular Biology: A Practical Approach", IRL Press at Oxford University Press, Oxford.
- A functional equivalent of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15 or SEQ ID NO:17 can be furthermore defined by the degree of homology or identity with SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15 or SEQ ID NO:17, respectively, and can furthermore comprise also natural or artificial mutations of the aforementioned nucleic acid sequences which encode a polypeptide with the activity of a nuclear encoded Clp-protease.
- The present invention also encompasses, for example, those nucleotide sequences which are obtained by modification of the SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15 or SEQ ID NO:17.
- For example, such modifications can be generated by techniques with which the skilled worker is familiar, such as "Site Directed Mutagenesis", "Error Prone PCR", "DNA-shuffling" (Nature 370, 1994, pp.389-391) or "Staggered Extension Process" (Nature Biotechnol. 16, 1998, pp.258-261). The aim of such a modification can be, for example, the insertion of further cleavage sites for restriction enzymes, the removal of DNA in order to truncate the sequence, the substitution of nucleotides to optimize the codons, or the addition of further sequences. Proteins which are encoded via modified nucleic acid sequences must retain the desired function despite a deviating nucleic acid sequence.

The term "functional equivalents" can also relate to the amino acid sequence encoded by the nucleic acid sequence in question. In this case, the term "functional equivalent" describes a protein whose amino acid sequence has a defined percentage of identity or
5 homology with SEQ ID NO:3.

Functional equivalents thus also comprise naturally occurring variants of the herein-described sequences and artificial nucleic acid sequences, for example those which have been obtained by chemical synthesis and which are adapted to the codon usage,
10 and also the amino acid sequences derived from them.

"Genetic control sequence" describes sequences which have an effect on the transcription and, if appropriate, translation of the nucleic acids according to the invention in prokaryotic or eukaryotic organisms. Examples thereof are promoters, terminators or
15 what are known as "enhancer" sequences. In addition to these control sequences, or instead of these sequences, the natural regulation of these sequences may still be present before the actual structural genes and may, if appropriate, have been genetically modified in such a way that the natural regulation has been switched off and the expression of the target gene has been modified, that is to say increased or reduced. The
20 choice of the control sequence depends on the host organism or starting organism. Genetic control sequences furthermore also comprise the 5'-untranslated region, introns or the noncoding 3'-region of genes. Control sequences are furthermore understood as meaning those which make possible homologous recombination or insertion into the genome of a host organism or which permit removal from the genome. Genetic
25 control sequences also comprise further promoters, promoter elements or minimal promoters, and sequences which have an effect on the chromatin structure (for example matrix attachment regions (MARs)), which can modify the expression-governing properties. Thus, genetic control sequences may bring about for example the additional dependence of the tissue-specific expression on certain stress factors. Such elements
30 have been described, for example, for water stress, abscisic acid (Lam E and Chua NH, J Biol Chem 1991; 266(26): 17131 -17135), high- and low-temperature stress (Plant Cell 1994, (6): 251-264) and heat stress (Molecular & General Genetics, 1989, 217(2-3): 246-53).

35 "Homology" between two nucleic acid sequences or polypeptide sequences is defined by the identity of the nucleic acid sequence/polypeptide sequence over in each case the entire sequence length, which is calculated by alignment with the aid of the program algorithm GAP according to Needleman and Wunsch 1970, J. Mol. Biol. 48; 443-453) setting the following parameters for polypeptides:

40

Gap Weight: 8 Length Weight: 2
Average Match: 2,912 Average Mismatch:-2,003

and the following parameters for nucleic acids:

Gap Weight: 50 Length Weight: 3
5 Average Match: 10.000 Average Mismatch: 0.000

In the following text, the term identity is also used synonymously with the term "homology".

10 "Mutations" of nucleic or amino acid sequences comprise substitutions, additions, deletions, inversions or insertions of one or more nucleotide residues, which may also bring about changes in the corresponding amino acid sequence of the target protein by substitution, insertion or deletion of one or more amino acids, although the functional properties of the target proteins are, overall, essentially retained.

15 "Natural genetic environment" means the natural chromosomal locus in the organism of origin. In the case of a genomic library, the natural genetic environment of the nucleic acid sequence is preferably retained at least in part. The environment flanks the nucleic acid sequence at least at the 5'- or 3'-side and has a sequence length of at least 50 bp,
20 preferably at least 100 bp, especially preferably at least 500 bp, very especially preferably at least 1000 bp, and most preferably at least 5000 bp.

25 "Plants" for the purposes of the invention are plant cells, plant tissues, plant organs, or intact plants, such as seeds, tubers, flowers, pollen, fruits, seedlings, roots, leaves, stems or other plant parts. Moreover, the term plants is understood as meaning propagation material such as seeds, fruits, seedlings, slips, tubers, cuttings or root stocks.

30 "Recombinant DNA" describes a combination of DNA sequences which can be generated by recombinant DNA technology.

35 "Recombinant DNA technology": generally known techniques for fusing DNA sequences (for example described in Sambrook et al., 1989, Cold Spring Harbor, NY, Cold Spring Harbor Laboratory Press).

40 "Replication origins" ensure the multiplication of the expression cassettes or vectors according to the invention in microorganisms and yeasts, for example the pBR322 ori or the P15A ori in *E. coli* (Sambrook et al.: "Molecular Cloning. A Laboratory Manual", 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989) and the ARS1 ori in yeast (Nucleic Acids Research, 2000, 28(10): 2060-2068).

45 "Reporter genes" encode readily quantifiable proteins. The transformation efficacy or the expression site or timing can be assessed by means of these genes via growth

assay, fluorescence assay, chemoluminescence assay, bioluminescence assay or resistance assay or via a photometric measurement (intrinsic color) or enzyme activity. Very especially preferred in this context are reporter proteins (Schenborn E, Groskreutz D. Mol Biotechnol. 1999; 13(1):29-44) such as the "green fluorescent protein" (GFP) (Gerdes HH and Kaether C, FEBS Lett. 1996; 389(1):44-47; Chui WL et al., Curr Biol 1996, 6:325-330; Leffel SM et al., Biotechniques. 23(5):912-8, 1997), chloramphenicol acetyl transferase, a luciferase (Giacomin, Plant Sci 1996, 116:59-72; Scikantha, J Bact 1996, 178:121; Millar et al., Plant Mol Biol Rep 1992 10:324-414), and luciferase genes, in general \square -galactosidase or \square -glucuronidase (Jefferson et al., EMBO J. 1987, 6, 3901-3907) or the Ura3 gene.

"Selection markers" confer resistance to antibiotics or other toxic compounds: examples which may be mentioned in this context are the neomycin phosphotransferase gene, which confers resistance to the aminoglycoside antibiotics neomycin (G 418), kanamycin, paromycin (Deshayes A et al., EMBO J. 4 (1985) 2731-2737), the sul gene, which encodes a mutated dihydropteroate synthase (Guerineau F et al., Plant Mol Biol. 1990; 15(1):127-136), the hygromycin B phosphotransferase gene (Gen Bank Accession NO: K 01193) and the shble resistance gene, which confers resistance to the bleomycin antibiotics such as zeocin. Further examples of selection marker genes are genes which confer resistance to 2-deoxyglucose-6-phosphate (WO 98/45456) or phosphinothricin and the like, or those which confer a resistance to antimetabolites, for example the dhfr gene (Reiss, Plant Physiol. (Life Sci. Adv.) 13 (1994) 142-149). Examples of other genes which are suitable are trpB or hisD (Hartman SC and Mulligan RC, Proc Natl Acad Sci U S A. 85 (1988) 8047-8051). Another suitable gene is the mannose phosphate isomerase gene (WO 94/20627), the ODC (ornithine decarboxylase) gene (McConlogue, 1987 in: Current Communications in Molecular Biology, Cold Spring Harbor Laboratory, Ed.) or the Aspergillus terreus deaminase (Tamura K et al., Biosci Biotechnol Biochem. 59 (1995) 2336-2338).

"Transformation" describes a process for introducing heterologous DNA into a pro- or eukaryotic cell. The term transformed cell describes not only the product of the transformation process per se, but also all of the transgenic progeny of the transgenic organism generated by the transformation.

"Target/target protein": a polypeptide encoded via the nucleic acid sequence according to the invention (this term is defined herein below), which may take the form of an enzyme in the traditional sense or, for example, of a structural protein, a protein relevant for developmental processes, regulatory protein such as transcription factors, kinases, phosphatases, receptors, channel subunits, transport proteins, regulatory subunits which confer substrate or activity regulation to an enzyme complex. All of the targets or sites of action share the characteristic that their functional presence is essential for survival or normal development and growth.

5 "Transgenic": referring to a nucleic acid sequence, an expression cassette or a vector comprising a nucleic acid sequence according to the invention or an organism transformed with the abovementioned nucleic acid sequence, expression cassette or vector, the term transgenic describes all those constructs which have been generated by genetic engineering methods in which either the nucleic acid sequence of the target protein or a genetic control sequence linked operably to the nucleic acid sequence of the target protein or a combination of the abovementioned possibilities are not in their natural genetic environment or have been modified by recombinant methods. In this context, the modification can be achieved, for example, by mutating one or more nucleotide residues of the nucleic acid sequence in question.

15 Intracellular protein degradation and its regulation are important for several processes like recycling of aminoacids, prevention of protein agglomeration and regulation of signaling processes (e.g. signalling of phytohormones). Cytosolic Proteins that are to be degraded are ubiquitinated at the N-terminus and delivered to the proteasome which is established as a complex of a large number of protein components in eucaryotes. Roughly 12% of the genes in *Arabidopsis thaliana* are encoding proteins involved in protein degradation by the ubiquitin pathway.

20 The stroma of plant chloroplasts contains a unique ubiquitin-independent, ATP-dependent protease consisting of two mayor components, a serine-type protease (ClpP) and an ATPase (ClpC, -D, -X) both of which are encoded by enzyme families in *Arabidopsis thaliana* (for details on the differing nomenclatures in literature see Adam et al. 2001, *Plant Physiology* 125, pp.1912-18). Six unique ClpP Isoforms (ClpP1-6) are nuclear encoded in *Arabidopsis* and at least one ClpP ist encoded in the plastid genome (pClpP) all of which carry the three conserved active site aminoacids characteristic for a catalytic triade of serine proteases. Further three nuclear encoded ClpP-Isoforms which miss the conserved amino acid residues of the catalytic triade are found in *Arabidopsis* (ClpR1, ClpR3, ClpR4). The catalytic activity of ClpR-type ClpP-Isoforms has not been shown so far.

35 At least one ClpP and two ClpX proteins may be targeted to mitochondria in *Arabidopsis* as deduced from N-terminal signal sequences. ClpP Proteases are conserved in bacteria. The ClpP protease in *E.coli* was formerly known as "protease Ti". A knock out of the protease Ti was shown to be not lethal. *E.coli* ClpP is assambled as a complex of 14 ClpP subunits in two heptameric rings. Co-immunoprecipitation suggests complexes of similar sizes and an ATP-dependet interaction of ClpP and ClpC subunits in Chloroplasts of *Arabidopsis thaliana* (Halperin et al. 2001, *Planta* 213, pp. 614-619). Furthermore, a 350kDa ClpP complex has been identified in *Arabidopsis* chloroplasts using blue native gel electrophoresis. The complex presumingly containins most of the known ClpP Isoenzymes (Benoit-Pellier et al. 2001, *Journal of Biological Chemistry*

276, pp. 16348-16327). Consequently the complexity and redundancy of plant Clp proteases is high and detailed information about composition of the clp complex and the functional role of its subunits remain to be clarified. Particularly the role of ClpP redundancy is still unclear.

5

The ClpP subunit is capable of actively hydrolysing peptides of max. 5 aminoacids in vitro. ClpA,B,C subunits constitute ATP-hydrolysing chaperones which unfold target-proteins and present them for hydrolysis to ClpP (Porankiewicz et al. 1999, Molecular Microbiology 32, 449-458). Involvement of ClpP in the degradation of the cytochrome b6f complex an PSII has been described in Chlamydomonas (Majeran et al. 2001, Plant Physiology 23+, pp. 421-433). Functional properties of ClpR-type Clp-Proteases as well as the ClpP like Proteases are yet to be determined.

Surprisingly, it has been found within the scope of the present invention that plants in which a Clp protease was reduced in a selective manner have phenotypes which are comparable with phenotypes generated by herbicide application. Drastic growth retardation and damage such as were observed.

The present invention relates to the use of a polypeptide, which has the activity of nuclear encoded Clp-protease in a method for identifying herbicides, preferably of a polypeptide, which has the activity of nuclear encoded Clp-protease, which is

- a) selected from the group consisting of ClpP1-protease, ClpP2-protease, ClpP3-protease, ClpP4-protease and ClpP6-protease; or
- 25 b) selected from the group consisting of ClpR1-protease, ClpR3-protease, ClpR4-protease; or
- c) ClpP-like-protease, wherein more preferably
- 30 a) the ClpP1-protease is encoded by a nucleic acid sequence which comprises:
 - i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:1, or
 - 35 ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:2 by back translating, or
 - 40 iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:1 which has an identity with SEQ ID NO:1 of has at least 50%; or

10

- iv) a functional equivalent of the nucleic acid sequence shown in SEQ ID NO:1, which is encoded by an amino acid sequence that has at least an identity of 50% with the SEQ ID NO:2;
- 5 b) the ClpP2-protease encoded by a nucleic acid sequence which comprises:
- i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:3, or
- 10 ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:4 by back translating, or
- 15 iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:3 which has an identity with SEQ ID NO:3 of has at least 50%; or
- iv) a functional equivalent of the nucleic acid sequence shown in SEQ ID NO:3, which is encoded by an amino acid sequence that has at least an identity of 50% with the SEQ ID NO:4;
- 20 c) the ClpP3-protease is encoded by a nucleic acid sequence which comprises:
- i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:5, or
- 25 ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:6 by back translating, or
- 30 iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:5 which has an identity with SEQ ID NO:5 of has at least 50%; or
- iv) a functional equivalent of the nucleic acid sequence shown in SEQ ID NO:5, which is encoded by an amino acid sequence that has at least an identity of 50% with the SEQ ID NO:6;
- 35 d) the ClpP4-protease is encoded by a nucleic acid sequence which comprises:
- i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:7, or
- 40

11

- ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:8 by back translating, or
- 5 iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:7 which has an identity with SEQ ID NO:7 of has at least 50%; or
- 10 iv) a functional equivalent of the nucleic acid sequence shown in SEQ ID NO:7, which is encoded by an amino acid sequence that has at least an identity of 50% with the SEQ ID NO:8;
- e) the ClpP6-protease is encoded by a nucleic acid sequence which comprises:
- 15 i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:9, or
- 20 ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:10 by back translating, or
- 25 iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:9 which has an identity with SEQ ID NO:9 of has at least 50%; or
- iv) a functional equivalent of the nucleic acid sequence shown in SEQ ID NO:9, which is encoded by an amino acid sequence that has at least an identity of 50% with the SEQ ID NO:10;
- f) the ClpR1-protease is encoded by a nucleic acid sequence which comprises:
- 30 i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:11, or
- 35 ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:12 by back translating, or
- 40 iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:11 which has an identity with SEQ ID NO:11 of has at least 50%; or
- iv) a functional equivalent of the nucleic acid sequence shown in SEQ ID NO:11, which is encoded by an amino acid sequence that has at least an identity of 50% with the SEQ ID NO:12;

- g) the ClpR3-protease is encoded by a nucleic acid sequence which comprises:
- 5 i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:13, or
- 10 ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:14 by back translating, or
- 15 iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:13 which has an identity with SEQ ID NO:13 of has at least 50%; or
- iv) a functional equivalent of the nucleic acid sequence shown in SEQ ID NO:13, which is encoded by an amino acid sequence that has at least an identity of 50% with the SEQ ID NO:14;
- h) the ClpR4-protease is encoded by a nucleic acid sequence which comprises:
- 20 i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:15, or
- 25 ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:16 by back translating, or
- 30 iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:15 which has an identity with SEQ ID NO:15 of has at least 50%; or
- iv) a functional equivalent of the nucleic acid sequence shown in SEQ ID NO:15, which is encoded by an amino acid sequence that has at least an identity of 50% with the SEQ ID NO:16;
- i) the ClpP like-protease is encoded by a nucleic acid sequence which comprises:
- 35 i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:17, or
- 40 ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:18 by back translating, or

13

iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:17 which has an identity with SEQ ID NO:17 of has at least 50%; or

5 iv) a functional equivalent of the nucleic acid sequence shown in SEQ ID NO:17, which is encoded by an amino acid sequence that has at least an identity of 50% with the SEQ ID NO:18;

wherein the sequences b) i-iv, e) i-iv, f) i-iv and are especially preferred.

10 The term "comprising" in relation to a nucleic acid sequence means that the nucleic acid sequence can be flanked by additional nucleic acid sequences that have on the 5' end and on the 3' end or on the 5' end or on the 3' end on the end a sequence length of at least 1000 bp, preferably at least 500 bp, more preferably at least 250bp, most preferably at least 100bp.

15 The functional equivalent according to the invention of SEQ ID NO:1 as described in a) iii), which encodes a polypeptide, which has the activity of nuclear encoded Clp-protease, and has at least an identity of 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64% or 65% or preferably of 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78% or 79% more preferably of 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90% most preferably of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% with SEQ ID NO:1.

20 The functional equivalents of the nucleic acid sequence SEQ ID NO:1 set forth in a) iv. are encoded by an amino acid sequence, which has the activity of nuclear encoded Clp-protease and has at least an identity of 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64% or 65% or preferably of 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78% or 79% more preferably of 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90% most preferably of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% with SEQ ID NO:2.

30 The functional equivalent according to the invention of SEQ ID NO:3 as described in b) iii), which encodes a polypeptide, which has the activity of nuclear encoded Clp-protease, and has at least an identity of 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64% or 65% or preferably of 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78% or 79% more preferably of 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90% most preferably of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% with SEQ ID NO:3.

40 An example of a functional equivalent of SEQ ID NO: 3 is the nucleic acid sequence of *Arabidopsis thaliana* (Gene Bank Acc. No. AB022327). This sequence is herein incorporated by reference.

14

The functional equivalents of the nucleic acid sequence set forth SEQ ID NO:3 in b) iv. are encoded by an amino acid sequence, which has the activity of nuclear encoded Clp-protease and has at least an identity of 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64% or 65% or preferably of 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78% or 79% more preferably of 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90% most preferably of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% with SEQ ID NO:4.

The functional equivalent according to the invention of SEQ ID NO:5 as described in c) iii), which encodes a polypeptide, which has the activity of nuclear encoded Clp-protease, and has at least an identity of 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64% or 65% or preferably of 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78% or 79% more preferably of 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90% most preferably of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% with SEQ ID NO:5.

The functional equivalents of the nucleic acid sequence SEQ ID NO:5 set forth in c) iv. are encoded by an amino acid sequence, which has the activity of nuclear encoded Clp-protease and has at least an identity of 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64% or 65% or preferably of 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78% or 79% more preferably of 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90% most preferably of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% with SEQ ID NO:6.

The functional equivalent according to the invention of SEQ ID NO:7 as described in d) iii), which encodes a polypeptide, which has the activity of nuclear encoded Clp-protease, and has at least an identity of 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64% or 65% or preferably of 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78% or 79% more preferably of 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90% most preferably of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% with SEQ ID NO:7.

The functional equivalents of the nucleic acid sequence set forth SEQ ID NO:7 in d) iv. are encoded by an amino acid sequence, which has the activity of nuclear encoded Clp-protease and has at least an identity of 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64% or 65% or preferably of 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78% or 79% more preferably of 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90% most preferably of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% with SEQ ID NO:7.

The functional equivalent according to the invention of SEQ ID NO:9 as described in e) iii), which encodes a polypeptide, which has the activity of nuclear encoded Clp-

15

- protease, and has at least an identity of 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64% or 65% or preferably of 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78% or 79% more preferably of 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90% most preferably of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% with SEQ ID NO:9.

- The functional equivalents of the nucleic acid sequence set forth SEQ ID NO:9 in e) iv. are encoded by an amino acid sequence, which has the activity of nuclear encoded Clp-protease and has at least an identity of 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64% or 65% or preferably of 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78% or 79% more preferably of 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90% most preferably of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% with SEQ ID NO:10.
- 15 The functional equivalent according to the invention of SEQ ID NO:11 as described in f) iii), which encodes a polypeptide, which has the activity of nuclear encoded Clp-protease, and has at least an identity of 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64% or 65% or preferably of 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78% or 79% more preferably of 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90% most preferably of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% with SEQ ID NO:11.

- The functional equivalents of the nucleic acid sequence SEQ ID NO:11 set forth in f) iv. are encoded by an amino acid sequence, which has the activity of nuclear encoded Clp-protease and has at least an identity of 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64% or 65% or preferably of 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78% or 79% more preferably of 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90% most preferably of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% with SEQ ID NO:12.

30 An example of a functional equivalent of SEQ ID NO: 11 is the nucleic acid sequence of *Arabidopsis thaliana* (Gene Bank Acc. No. AB022330). This sequence is herein incorporated by reference.

- 35 The functional equivalent according to the invention of SEQ ID NO:13 as described in g) iii), which encodes a polypeptide, which has the activity of nuclear encoded Clp-protease, and has at least an identity of 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64% or 65% or preferably of 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78% or 79% more preferably of 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90% most preferably of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% with SEQ ID NO:13.

16

The functional equivalents of the nucleic acid sequence SEQ ID NO:13 set forth in g) iv. are encoded by an amino acid sequence, which has the activity of nuclear encoded Clp-protease and has at least an identity of 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64% or 65% or preferably of 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78% or 79% more preferably of 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90% most preferably of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% with SEQ ID NO:14.

The functional equivalent according to the invention of SEQ ID NO:15 as described in h) iii), which encodes a polypeptide, which has the activity of nuclear encoded Clp-protease, and has at least an identity of 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64% or 65% or preferably of 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78% or 79% more preferably of 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90% most preferably of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% with SEQ ID NO:15.

The functional equivalents of the nucleic acid sequence SEQ ID NO:15 set forth in h) iv. are encoded by an amino acid sequence, which has the activity of nuclear encoded Clp-protease and has at least an identity of 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64% or 65% or preferably of 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78% or 79% more preferably of 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90% most preferably of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% with SEQ ID NO:16.

The functional equivalent according to the invention of SEQ ID NO:17 as described in i) iii), which encodes a polypeptide, which has the activity of nuclear encoded Clp-protease, and has at least an identity of 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64% or 65% or preferably of 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78% or 79% more preferably of 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90% most preferably of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% with SEQ ID NO:17.

The functional equivalents of the nucleic acid sequence SEQ ID NO:17 set forth in i) iv. are encoded by an amino acid sequence, which has the activity of nuclear encoded Clp-protease and has at least an identity of 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64% or 65% or preferably of 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78% or 79% more preferably of 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89% or 90% most preferably of 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% with SEQ ID NO:18.

40

17

An example of a functional equivalent of SEQ ID NO: 17 is the nucleic acid sequence of *Arabidopsis thaliana* (Gene Bank Acc. No. AK118525). This sequence is herein incorporated by reference.

- 5 Furthermore claimed within the scope of the present invention are plant nucleic acid sequence

I) encoding a ClpP2-protease comprising:

- 10 a) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:3, or
- b) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID
- 15 NO:4 by backtranslating, or
- c) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:1 which has an identity with SEQ ID NO:3 of has at least 66%; or
- 20 d) a functional equivalent of the nucleic acid sequence shown in SEQ ID NO:11, which is encoded by an amino acid sequence that has at least an identity of 76% with the SEQ ID NO:4;

II) encoding a ClpR1-protease comprising:

- 25 a) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:11, or
- b) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID
- 30 NO:12 by backtranslating, or
- c) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:1 which has an identity with SEQ ID NO:11 of has at least 69%; or
- 35 d) a functional equivalent of the nucleic acid sequence shown in SEQ ID NO:11, which is encoded by an amino acid sequence that has at least an identity of 71% with the SEQ ID NO:12;

40 III) encoding a ClpP-like-protease comprising:

- a) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:17, or
- 5 b) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:18 by backtranslating, or
- c) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:1 which has an identity with SEQ ID NO:17 of has at least 67%; or
- 10 d) a functional equivalent of the nucleic acid sequence shown in SEQ ID NO:17, which is encoded by an amino acid sequence that has at least an identity of 79% with the SEQ ID NO:18;

15 The functional equivalent of SEQ ID NO:3 set forth in I c) has at least an identity of 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, by preference at least 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82% or 83%, preferably at least 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92% or 93%, especially preferably at least 94%, 95%, 96%, 97%, 98% or 99% with SEQ ID NO:3.

20 The functional equivalents of the nucleic acid sequence SEQ ID NO:3 set forth in I) d) are encoded by an amino acid sequence, which has the activity of nuclear encoded Clp-protease and has at least an identity of 77%, by preference at least 78%, 79%, 80%, 81%, 82% or 83%, preferably at least 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, especially preferably at least 94%, 95%, 96%, 97%, 98%, 99% with

25 SEQ ID NO:4.

The functional equivalent of SEQ ID NO:11 set forth in II c) has at least an identity of 69%, 70%, 71%, 72%, 73% or 74%, by preference at least 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82% or 83%, preferably at least 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92% or 93%, especially preferably at least 94%, 95%, 96%, 97%, 98% or 99% with SEQ ID NO:11.

30

The functional equivalents of the nucleic acid sequence SEQ ID NO:11 set forth in II) d) are encoded by an amino acid sequence, which has the activity of nuclear encoded Clp-protease and has at least an identity of 71% by preference at least 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, preferably at least 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, especially preferably at least 94%, 95%, 96%, 97%, 98%, 99% with SEQ ID NO:12.

35

40 The functional equivalent of SEQ ID NO:17 set forth in I c) has at least an identity of 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, by preference at least 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82% or 83%, preferably at least 84%, 85%, 86%, 87%, 88%,

89%, 90%, 91%, 92% or 93%, especially preferably at least 94%, 95%, 96%, 97%, 98% or 99% with SEQ ID NO: 17.

5 The functional equivalents of the nucleic acid sequence SEQ ID NO:17 set forth in I) d) are encoded by an amino acid sequence, which has the activity of nuclear encoded Clp-protease and has at least an identity of 79%, by preference at least 79%, 80%, 81%, 82% or 83%, preferably at least 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, especially preferably at least 94%, 95%, 96%, 97%, 98%, 99% with SEQ ID NO:18.

10

The polypeptides encoded by the abovementioned nucleic acid sequences according to I c)-d), II c)-d) and III c)-d) are likewise claimed. The functional equivalents as described in c) and d) are distinguished by the same functionality, i.e. they have the activity of a clp-protease.

15

The nucleic acid sequences I c)-d), II c)-d) and III c)-d) are hereinbelow termed NCLP-sequences.

20 The term "nucleic acid sequences according to the invention" which is used hereinbelow refers to nucleic acid sequences encoding a polypeptide, which has the activity of nuclear encoded Clp-protease in a method for identifying herbicides, preferably of a polypeptide, which has the activity of nuclear encoded Clp-protease, which is

- 25 a) selected from the group consisting of ClpP1-protease, ClpP2-protease, ClpP3-protease, ClpP4-protease and ClpP6-protease; or
- b) selected from the group consisting of ClpR1-protease, ClpR3-protease, ClpR4-protease; or

30 c) ClpP-like-protease, wherein more preferably

- a) the ClpP1-protease is encoded by a nucleic acid sequence which comprises:
- 35 i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:1, or
- ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:2 by back translating, or
- 40 iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:1 which has an identity with SEQ ID NO:1 of has at least 50%; or

- 5 iv) a functional equivalent of the nucleic acid sequence shown in SEQ ID NO:1, which is encoded by an amino acid sequence that has at least an identity of 50% with the SEQ ID NO:2;
- 10 b) the ClpP2-protease encoded by a nucleic acid sequence which comprises:
- i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:3, or
- ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:4 by back translating, or
- 15 iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:3 which has an identity with SEQ ID NO:3 of has at least 50%; or
- 20 iv) a functional equivalent of the nucleic acid sequence shown in SEQ ID NO:3, which is encoded by an amino acid sequence that has at least an identity of 50% with the SEQ ID NO:4;
- c) the ClpP3-protease is encoded by a nucleic acid sequence which comprises:
- 25 i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:5, or
- ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:6 by back translating, or
- 30 iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:5 which has an identity with SEQ ID NO:5 of has at least 50%; or
- 35 iv) a functional equivalent of the nucleic acid sequence shown in SEQ ID NO:5, which is encoded by an amino acid sequence that has at least an identity of 50% with the SEQ ID NO:6;
- d) the ClpP4-protease is encoded by a nucleic acid sequence which comprises:
- 40 i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:7, or

21

- ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:8 by back translating, or
- 5 iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:7 which has an identity with SEQ ID NO:7 of has at least 50%; or
- 10 iv) a functional equivalent of the nucleic acid sequence shown in SEQ ID NO:7, which is encoded by an amino acid sequence that has at least an identity of 50% with the SEQ ID NO:8;
- e) the ClpP6-protease is encoded by a nucleic acid sequence which comprises:
- 15 i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:9, or
- 20 ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:10 by back translating, or
- 25 iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:9 which has an identity with SEQ ID NO:9 of has at least 50%; or
- iv) a functional equivalent of the nucleic acid sequence shown in SEQ ID NO:9, which is encoded by an amino acid sequence that has at least an identity of 50% with the SEQ ID NO:10;
- f) the ClpR1-protease is encoded by a nucleic acid sequence which comprises:
- 30 i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:11, or
- 35 ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:12 by back translating, or
- 40 iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:11 which has an identity with SEQ ID NO:11 of has at least 50%; or
- iv) a functional equivalent of the nucleic acid sequence shown in SEQ ID NO:11, which is encoded by an amino acid sequence that has at least an identity of 50% with the SEQ ID NO:12;

- g) the ClpR3-protease is encoded by a nucleic acid sequence which comprises:
- 5 i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:13, or
- ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:14 by back translating, or
- 10 iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:13 which has an identity with SEQ ID NO:13 of has at least 50%; or
- iv) a functional equivalent of the nucleic acid sequence shown in SEQ ID NO:13, which is encoded by an amino acid sequence that has at least an identity of 50% with the SEQ ID NO:14;
- 15
- h) the ClpR4-protease is encoded by a nucleic acid sequence which comprises:
- 20 i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:15, or
- ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:16 by back translating, or
- 25 iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:15 which has an identity with SEQ ID NO:15 of has at least 50%; or
- iv) a functional equivalent of the nucleic acid sequence shown in SEQ ID NO:15, which is encoded by an amino acid sequence that has at least an identity of 50% with the SEQ ID NO:16;
- 30
- i) the ClpP like-protease is encoded by a nucleic acid sequence which comprises:
- 35 i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:17, or
- ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:18 by back translating, or
- 40

23

- iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:17 which has an identity with SEQ ID NO:17 of has at least 50%; or
- iv) a functional equivalent of the nucleic acid sequence shown in SEQ ID NO:17, which is encoded by an amino acid sequence that has at least an identity of 50% with the SEQ ID NO:18;

wherein the sequences b) i-iv, e) i-iv, f) i-iv and are especially preferred

- A polypeptide, which has the activity of nuclear encoded Clp-protease and is encoded by a nucleic acid sequence according to the invention are hereinbelow simply referred to as "CLP".

- Reduced amounts of glyoxysomal CLP cause growth retardation and necrotic and chlorotic leaves in plants.

- The gene products of the nucleic acids according to the invention constitute novel targets for herbicides, which make possible the provision of novel herbicides for controlling undesired plants. Moreover, the gene products of the nucleic acids according to the invention constitute novel targets for growth regulators which make possible the provision of novel growth regulators for regulating the growth of plants.

- Undesired plants are understood as meaning, in the broadest sense, all those plants which grow at locations where they are undesired, for example:

- Dicotyledonous weeds of the genera: Sinapis, Lepidium, Galium, Stellaria, Matricaria, Anthemis, Galinsoga, Chenopodium, Urtica, Senecio, Amaranthus, Portulaca, Xanthium, Convolvulus, Ipomoea, Polygonum, Sesbania, Ambrosia, Cirsium, Carduus, Sonchus, Solanum, Rorippa, Rotala, Lindernia, Lamium, Veronica, Abutilon, Emex, Datura, Viola, Galeopsis, Papaver, Centaurea, Trifolium, Ranunculus, Taraxacum.

- Monocotyledonous weeds from the genera: Echinochloa, Setaria, Panicum, Digitaria, Phleum, Poa, Festuca, Eleusine, Brachiaria, Lolium, Bromus, Avena, Cyperus, Sorghum, Agropyron, Cynodon, Monochoria, Fimbristylis, Sagittaria, Eleocharis, Scirpus, Paspalum, Ischaemum, Sphenoclea, Dactyloctenium, Agrostis, Alopecurus, Apera.

- SEQ ID NO:1; 3, 5, 7, 9, 11, 13, 15, 17 or 19-21 or parts of SEQ ID NO: 1; 3, 5, 7, 9, 11, 13, 15, 17 or 19-21 can be used for the preparation of hybridization probes. The preparation of these probes and the experimental procedure is known. For example, this can be effected via the selective preparation of radioactive or nonradioactive probes by PCR and the use of suitably labeled oligonucleotides, followed by hybridization experiments. The technologies required for this purpose are detailed, for example,

in T. Maniatis, E.F. Fritsch and J. Sambrook, "Molecular Cloning: A Laboratory Manual", Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1989). The probes in question can furthermore be modified by standard technologies (Lit. SDM or random mutagenesis) in such a way that they can be employed for further purposes, for example as a probe which hybridizes specifically with mRNA and the corresponding coding sequences in order to analyze the corresponding sequences in other organisms.

The abovementioned probes can be used for the detection and isolation of functional equivalents of SEQ ID NO:2, 4, 6, 8, 10, 12, 14, 16 or 18 from other plant species on the basis of sequence identities. In this context, part or all of the sequence of the SEQ ID NO:2 in question is used as a probe for screening a genomic or cDNA library of the plant species in question or in a computer search for sequences of functional equivalents in electronic databases.

Preferred plant species are the undesired plants which have already been mentioned at the outset.

The invention furthermore relates to expression cassettes comprising

- a) genetic control sequences in operable linkage with a NCLP sequence; or
- b) additional functional elements, or
- c) a combination of a) and b);

and to the use of expression cassettes comprising

- a) genetic control sequences in operable linkage with a nucleic acid sequence according to the invention,
- b) additional functional elements, or
- c) a combination of a) and b);

for expressing a CLP, which can be used in in vitro assay systems. Both embodiments of the above-described expression cassettes are referred in the following text as expression cassette according to the invention.

In a preferred embodiment, an expression cassette according to the invention comprises a promoter at the 5' end of the coding sequence and, at the 3' end, a transcription termination signal and, if appropriate, further genetic control sequences which are linked operably with the interposed nucleic acid sequence according to the invention.

The expression cassettes according to the invention are also understood as meaning analogs which can be brought about, for example, by a combination of the individual nucleic acid sequences on a polynucleotide (multiple constructs), on a plurality of
5 polynucleotides in a cell (cotransformation) or by sequential transformation.

Advantageous genetic control sequences under point a) for the expression cassettes according to the invention or for vectors comprising expression cassettes according to the invention are, for example, promoters such as the cos, tac, trp, tet, lpp, lac, lacIq,
10 T7, T5, T3, gal, trc, ara, SP6, \square -PR or the \square -PL promoter, all of which can be used for expressing a CLP, in Gram-negative bacterial strains.

Examples of further advantageous genetic control sequences are present, for example, in the promoters amy and SPO2, both of which can be used for expressing a CLP, in
15 Gram-positive bacterial strains, and in the yeast or fungal promoters AUG1, GPD-1, PX6, TEF, CUP1, PGK, GAP1, TPI, PHO5, AOX1, GAL10/CYC1, CYC1, OliC, ADH, TDH, Kex2, MFA or NMT or combinations of the abovementioned promoters (Degryse et al., Yeast 1995 June 15; 11(7):629-40; Romanos et al. Yeast 1992 June;8(6):423-88; Benito et al. Eur. J. Plant Pathol. 104, 207-220 (1998); Cregg et al. Biotechnology (N Y) 1993 Aug;11(8):905-10; Luo X., Gene 1995 Sep 22;163(1):127-31; Nacken et al.,
20 Gene 1996 Oct 10;175(1-2): 253-60; Turgeon et al., Mol Cell Biol 1987 Sep;7(9):3297-305) or the transcription terminators NMT, Gcy1, TrpC, AOX1, nos, PGK or CYC1 (Degryse et al., Yeast 1995 June 15; 11(7):629-40; Brunelli et al. Yeast 1993 (Dec9(12): 1309-18; Frisch et al., Plant Mol. Biol. 27(2), 405-409 (1995); Scorer et al., Biotechnology (N.Y. 12 (2), 181-184 (1994), Genbank acc. number Z46232; Zhao et al. Genbank acc number : AF049064; Punt et al., (1987) Gene 56 (1), 117-124), all of which can be
25 used for expressing CLP, in yeast strains.

Examples of genetic control sequences which are suitable for expression in insect cells are the polyhedrin promoter and the p10 promoter (Luckow, V.A. and Summers, M.D. (1988) Bio/Techn. 6, 47-55).
30

Advantageous genetic control sequences for expressing CLP, in cell culture, in addition to polyadenylation sequences such as, for example, from simian virus 40, are eukaryotic promoters of viral origin such as, for example, promoters of the polyoma virus, adenovirus 2, cytomegalovirus or simian virus 40.
35

Further advantageous genetic control sequences for expressing nuclear encoded Clp Protease, in plants are present in the plant promoters CaMV/35S [Franck et al., Cell 21(1980) 285-294], PRP1 [Ward et al., Plant. Mol. Biol. 22 (1993)], SSU, OCS, LEB4, USP, STLS1, B33, NOS; FBpaseP (WO 98/18940) or in the ubiquitin or phaseolin promoter; a promoter which is preferably used being, in particular, a plant promoter or
40

a promoter derived from a plant virus. Especially preferred are promoters of viral origin such as the promoter of the cauliflower mosaic virus 35S transcript (Franck et al., Cell 21 (1980), 285-294; Odell et al., Nature 313 (1985), 810-812). Further preferred constitutive promoters are, for example, the agrobacterium nopaline synthase promoter, the

5 TR double promoter, the agrobacterium OCS (octopine synthase) promoter, the ubiquitin promoter, (Holtorf S et al., Plant Mol Biol 1995, 29:637-649), the promoters of the vacuolar ATPase subunits, or the promoter of a proline-rich wheat protein (WO 91/13991).

- 10 The expression cassettes may also comprise, as genetic control sequence, a chemically inducible promoter, by which the expression of the exogenous gene in the plant can be controlled at a specific point in time. Such promoters, such as, for example, the PRP1 promoter (Ward et al., Plant. Mol. Biol. 22 (1993), 361-366), a salicylic-acid-inducible promoter (WO 95/19443), a benzenesulfonamide-inducible promoter (EP-A-
- 15 0388186), a tetracyclin-inducible promoter (Gatz et al., (1992) Plant J. 2, 397404), an abscisic-acid-inducible promoter (EP-A 335528) or an ethanol- or cyclohexanone-inducible promoter (WO 93/21334) may also be used.

- Furthermore, suitable promoters are those which confer tissue- or organ-specific expression in, for example, anthers, ovaries, flowers and floral organs, leaves, stomata, trichomes, stems, vascular tissues, roots and seeds. Others which are suitable in addition to the abovementioned constitutive promoters are, in particular, those promoters which ensure leaf-specific expression. Promoters which must be mentioned are the
- 20 potato cytosolic FBPase promoter (WO 97/05900), the rubisco (ribulose-1,5-bisphosphate carboxylase) SSU (small subunit) promoter or the ST-LSI promoter from potato (Stockhaus et al., EMBO J. 8 (1989), 2445 - 245). Promoters which are furthermore preferred are those which control expression in seeds and plant embryos. Examples of seed-specific promoters are the phaseolin promoter (US 5,504,200, Bustos MM et al., Plant Cell. 1989;1(9):839-53), the promoter of the 2S albumin gene (Joseffson
- 25 LG et al., J Biol Chem 1987, 262:12196-12201), the legumin promoter (Shirsat A et al., Mol Gen Genet. 1989;215(2):326-331), the USP (unknown seed protein) promoter (Bäumlein H et al., Molecular & General Genetics 1991, 225(3):459-67), the napin gene promoter (Stalberg K, et al., L. Planta 1996, 199:515-519), the sucrose binding protein promoter (WO 00/26388) or the LeB4 promoter (Bäumlein H et al., Mol Gen
- 30 Genet 1991, 225: 121-128; Fiedler, U. et al., Biotechnology (NY) (1995), 13 (10) 1090).
- 35

- Further promoters which are suitable as genetic control sequences are, for example, specific promoters for tubers, storage roots or roots, such as, for example, the class I patatin promoter (B33), the potato cathepsin D inhibitor promoter, the starch synthase
- 40 (GBSS1) promoter or the sporamin promoter, fruit-specific promoters such as, for example, the fruit-specific promoter from tomato (EP-A 409625), fruit-maturation-specific promoters such as, for example, the fruit-maturation-specific promoter from tomato

(WO 94/21794), inflorescence-specific promoters such as, for example, the phytoene synthase promoter (WO 92/16635) or the promoter of the P- π gene (WO 98/22593), or plastid- or chromoplast-specific promoters such as, for example, the RNA polymerase promoter (WO 97/06250), or else the Glycine max phosphoribosyl-pyrophosphate amidotransferase promoter (see also Genbank Accession No. U87999), or another node-specific promoter as described in EP-A 249676.

Additional functional elements b) are understood as meaning, by way of example but not by limitation, reporter genes, replication origins, selection markers and what are known as affinity tags, in fusion with CLP, directly or by means of a linker optionally comprising a protease cleavage site. Further suitable additional functional elements are sequences which ensure that the product is targeted into the apoplasts, into plastids, the vacuole, the mitochondrion, the peroxisome, the endoplasmatic reticulum (ER) or, owing to the absence of such operative sequences, remains in the compartment where it is formed, the cytosol, (Kermode, Crit. Rev. Plant Sci. 15, 4 (1996), 285-423).

Also in accordance with the invention are vectors comprising at least one copy of the nucleic acid sequences according to the invention and/or the expression cassettes according to the invention.

In addition to plasmids, vectors are furthermore also understood as meaning all of the other known vectors with which the skilled worker is familiar, such as, for example, phages, viruses such as SV40, CMV, baculovirus, adenovirus, transposons, IS elements, phasmids, phagemids, cosmids or linear or circular DNA. These vectors can be replicated autonomously in the host organism or replicated chromosomally; chromosomal replication is preferred.

In a further embodiment of the vector, the nucleic acid construct according to the invention can advantageously also be introduced into the organisms in the form of a linear DNA and integrated into the genome of the host organism via heterologous or homologous recombination. This linear DNA may consist of a linearized plasmid or only of the nucleic acid construct as vector, or the nucleic acid sequences used.

Further prokaryotic or eukaryotic expression systems are mentioned in Chapters 16 and 17 in Sambrook et al., "Molecular Cloning: A Laboratory Manual." 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989. Further advantageous vectors are described in Hellens et al. (Trends in plant science, 5, 2000).

The expression cassette according to the invention and vectors derived therefrom can be used for transforming bacteria, cyanobacteria, (for example of the genus Synechocystis, Anabaena, Calothrix, Scytonema, Oscillatoria, Plectonema and Nostoc), proteo-

bacteria such as, for example, *Magnetococcus* sp. MC1, yeasts, filamentous fungi and algae and eukaryotic nonhuman cells (for example insect cells) with the aim of producing CLP, recombinantly, the generation of a suitable expression cassette depending on the organism in which the gene is to be expressed.

5

Vectors comprising a NCLP sequence form part of the subject-matter of the present invention.

10

In a further advantageous embodiment, the nucleic acid sequences according to the invention may also be introduced into an organism by themselves.

15

If, in addition to the nucleic acid sequences, further genes are to be introduced into the organism, they can all be introduced into the organism together in a single vector, or each individual gene can be introduced into the organism in each case in one vector, it being possible to introduce the different vectors simultaneously or in succession.

20

In this context, the introduction, into the organisms in question (transformation), of the nucleic acid(s) according to the invention, of the expression cassette or of the vector can be effected in principle by all methods with which the skilled worker is familiar.

25

30

In the case of microorganisms, the skilled worker will find suitable methods in the textbooks by Sambrook, J. et al. (1989) "Molecular cloning: A laboratory manual", Cold Spring Harbor Laboratory Press, von F.M. Ausubel et al. (1994) "Current protocols in molecular biology", John Wiley and Sons, by D.M. Glover et al., DNA Cloning Vol.1, (1995), IRL Press (ISBN 019-963476-9), by Kaiser et al. (1994) Methods in Yeast Genetics, Cold Spring Harbor Laboratory Press or Guthrie et al. "Guide to Yeast Genetics and Molecular Biology", Methods in Enzymology, 1994, Academic Press. In the transformation of filamentous fungi, the methods of choice are firstly the generation of protoplasts and transformation with the aid of PEG (Wiebe et al. (1997) Mycol. Res. 101 (7): 971-977; Proctor et al. (1997) Microbiol. 143, 2538-2591), and secondly the transformation with the aid of *Agrobacterium tumefaciens* (de Groot et al. (1998) Nat. Biotech. 16, 839-842).

35

40

In the case of dicots, the methods which have been described for the transformation and regeneration of plants from plant tissues or plant cells can be exploited for transient or stable transformation. Suitable methods are the biolistic method or the transformation of protoplasts (cf., for example, Willmitzer, L., 1993 Transgenic plants. In: Biotechnology, A Multi-Volume Comprehensive Treatise (H.J. Rehm, G. Reed, A. Pühler, P. Stadler, eds.), Vol. 2, 627-659, VCH Weinheim-New York-Basle-Cambridge), electroporation, the incubation of dry embryos in DNA-containing solution, microinjection and the agrobacterium-mediated gene transfer. The abovementioned methods are described, for example, in B. Jenet et al., Techniques for Gene Transfer,

in: Transgenic Plants, Vol. 1, Engineering and Utilization, edited by S.D. Kung and R. Wu, Academic Press (1993) 128-143 and in Potrykus, Annu. Rev. Plant Physiol. Plant Molec.Biol. 42 (1991) 205-225).

- 5 The transformation by means of agrobacteria, and the vectors to be used for the transformation, are known to the skilled worker and described extensively in the literature (Bevan et al., Nucl. Acids Res. 12 (1984) 8711. The intermediary vectors can be integrated into the agrobacterial Ti or Ri plasmid by means of homologous recombination owing to sequences which are homologous to sequences in the T-DNA. This plasmid
- 10 additionally contains the vir region, which is required for the transfer of the T-DNA. Intermediary vectors are not capable of replication in agrobacteria. The intermediary vector can be transferred to Agrobacterium tumefaciens by means of a helper plasmid (conjugation). Binary vectors are capable of replication both in E. coli and in agrobacteria. They contain a selection marker gene and a linker or polylinker which are framed
- 15 by the right and left T-DNA border region. They can be transformed directly into the agrobacteria (Holsters et al. Mol. Gen. Genet. 163 (1978), 181-187), EP A 0 120 516; Hoekema, in: The Binary Plant Vector System Offsetdrukkerij Kanters B.V., Alblasterdam (1985), Chapter V; Fraley et al., Crit. Rev. Plant. Sci., 4: 1-46 and An et al. EMBO J. 4 (1985), 277-287).
- 20 The transformation of monocots by means of vectors based on agrobacterium has also been described (Chan et al., Plant Mol. Biol. 22(1993), 491-506; Hiei et al., Plant J. 6 (1994) 271-282; Deng et al. Science in China 33 (1990), 28-34; Wilmink et al., Plant Cell Reports 11,(1992) 76-80; May et al. Biotechnology 13 (1995) 486-492; Conner and
- 25 Domisse; Int. J. Plant Sci. 153 (1992) 550-555; Ritchie et al. Transgenic Res. (1993) 252-265). Alternative systems for the transformation of monocots are the transformation by means of biolistic approach (Wan and Lemaux; Plant Physiol. 104 (1994), 37-48; Vasil et al. Biotechnology 11 (1992), 667-674; Ritala et al., Plant Mol. Biol 24, (1994) 317-325; Spencer et al., Theor. Appl. Genet. 79 (1990), 625-631), protoplast
- 30 transformation, the electroporation of partially permeabilized cells, and the introduction of DNA by means of glass fibers. In particular the transformation of maize has been described repeatedly in the literature (cf., for example, WO 95/06128; EP 0513849 A1; EP 0465875 A1; EP 0292435 A1; Fromm et al., Biotechnology 8 (1990), 833-844; Gordon-Kamm et al., Plant Cell 2 (1990), 603-618; Koziel et al., Biotechnology
- 35 11(1993) 194-200; Moroc et al., Theor Applied Genetics 80 (190) 721-726).

The successful transformation of other cereal species has also already been described for example in the case of barley (Wan and Lemaux, see above; Ritala et al., see above; wheat (Nehra et al., Plant J. 5(1994) 285-297).

Agrobacteria which have been transformed with a vector according to the invention can likewise be used in a known manner for the transformation of plants, such as test

30

plants like Arabidopsis or crop plants like cereals, maize, oats, rye, barley, wheat, soya, rice, cotton, sugarbeet, canola, sunflower, flax, hemp, potato, tobacco, tomato, carrot, capsicum, oilseed rape, tapioca, cassava, arrowroot, Tagetes, alfalfa, lettuce and the various tree, nut and grapevine species, for example by bathing scarified leaves or leaf segments in an agrobacterial solution and subsequently growing them in suitable media.

The genetically modified plant cells can be regenerated via all methods with which the skilled worker is familiar. Such methods can be found in the abovementioned publications by S.D. Kung and R. Wu, Potrykus or Höfgen and Willmitzer.

The transgenic organisms generated by transformation with one of the above-described embodiments of an expression cassette comprising a nucleic acid sequence according to the invention or a vector comprising the abovementioned expression cassette, and the recombinant CLP, which can be obtained from the transgenic organism by means of expression, form part of the subject matter of the present invention. The use of transgenic organisms comprising an expression cassette according to the invention, for example for providing recombinant protein, and/or the use of these organisms in in-vivo assay systems likewise form part of the subject matter of the present invention.

Preferred organisms for the recombinant expression are not only bacteria, yeasts, mosses, algae and fungi, but also eukaryotic cell lines.

Preferred mosses are Physcomitrella patens or other mosses described in Kryptogamen [Cryptogamia], Vol.2, Moose, Farne [Mosses, Ferns], 1991, Springer Verlag (ISBN 3540536515).

Preferred within the bacteria are, for example, bacteria from the genus Escherichia, Erwinia, Flavobacterium, Alcaligenes or cyanobacteria, for example from the genus Synechocystis, Anabaena, Calothrix, Scytonema, Oscillatoria, Plectonema and Nostoc, especially preferably Synechocystis or Anabaena.

Preferred yeasts are Candida, Saccharomyces, Schizosaccharomyces, Hansenula or Pichia.

Preferred fungi are Aspergillus, Trichoderma, Ashbya, Neurospora, Fusarium, Beauveria, Mortierella, Saprolegnia, Pythium, or other fungi described in Indian Chem Engr. Section B. Vol 37, No 1,2 (1995).

Preferred plants are selected in particular among monocotyledonous crop plants such as, for example, cereal species such as wheat, barley, sorghum or millet, rye, triticale,

- maize, rice or oats, and sugarcane. The transgenic plants according to the invention are, furthermore, in particular selected from among dicotyledonous crop plants such as, for example, Brassicaceae such as oilseed rape, cress, Arabidopsis, cabbages or canola; Leguminosae such as soyabean, alfalfa, pea, beans or peanut, Solanaceae such as potato, tobacco, tomato, egg plant or capsicum; Asteraceae such as sunflower, Tagetes, lettuce or Calendula; Cucurbitaceae such as melon, pumpkin/squash or zucchini, or linseed, cotton, hemp, flax, red pepper, carrot, sugar beet, or various tree, nut and grapevine species.
- 10 In principle, transgenic animals such as, for example, *C. elegans*, are also suitable as host organisms.
- Also preferred is the use of expression systems and vectors which are available to the public or commercially available.
- 15 Those which must be mentioned for use in *E. coli* bacteria are the typical advantageous commercially available fusion and expression vectors pGEX [Pharmacia Biotech Inc; Smith, D.B. and Johnson, K.S. (1988) *Gene* 67:31-40], pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ), which contains glutathione S transferase (GST), maltose binding protein or protein A, the pTrc vectors (Amann et al., (1988) *Gene* 69:301-315), "pKK233-2" from CLONTECH, Palo Alto, CA and the "pET", and the "pBAD" vector series from Stratagene, La Jolla and the TOPO-TA vector series from Invitrogen.
- 25 Further advantageous vectors for use in yeast are pYepSec1 (Baldari, et al., (1987) *Embo J.* 6:229-234), pMFa (Kurjan and Herskowitz, (1982) *Cell* 30:933-943), pJRY88 (Schultz et al., (1987) *Gene* 54:113-123), and pYES derivatives, pGAPZ derivatives, pPICZ derivatives, and the vectors of the "Pichia Expression Kit" (Invitrogen Corporation, San Diego, CA). Vectors for use in filamentous fungi are described in: van den Hondel, C.A.M.J.J. & Punt, P.J. (1991) "Gene transfer systems and vector development for filamentous fungi, in: *Applied Molecular Genetics of Fungi*, J.F. Peberdy, et al., eds., p. 1-28, Cambridge University Press: Cambridge.
- 30 As an alternative, insect cell expression vectors may also be used advantageously, for example for expression in Sf9, Sf21 or Hi5 cells, which are infected via recombinant Baculoviruses. Examples of these are the vectors of the pAc series (Smith et al. (1983) *Mol. Cell Biol.* 3:2156-2165) and the pVL series (Lucklow and Summers (1989) *Virology* 170:31-39). Others which may be mentioned are the Baculovirus expression systems "MaxBac 2.0 Kit" and "Insect Select System" from Invitrogen, Carlsbad or
- 40 "BacPAK Baculovirus Expression System" from CLONTECH, Palo Alto, CA. Insect cells are particularly suitable for overexpressing eukaryotic proteins since they effect posttranslational modifications of the proteins which are not possible in bacteria and

yeasts. The skilled worker is familiar with the handling of cultured insect cells and with their infection for expressing proteins, which can be carried out analogously to known methods (Luckow and Summers, Bio/Tech. 6, 1988, pp.47-55; Glover and Hames (eds) in DNA Cloning 2, A practical Approach, Expression Systems, Second Edition, 5 Oxford University Press, 1995, 205-244).

Plant cells or algal cells are others which can be used advantageously for expressing genes. Examples of plant expression vectors can be found as mentioned above in 10 Becker, D., et al. (1992) "New plant binary vectors with selectable markers located proximal to the left border", Plant Mol. Biol. 20: 1195-1197 or in Bevan, M.W. (1984) "Binary Agrobacterium vectors for plant transformation", Nucl. Acid. Res. 12: 8711-8721.

Moreover, the nucleic acid sequences according to the invention can be expressed in 15 mammalian cells. Examples of suitable expression vectors are pCDM8 and pMT2PC, which are mentioned in: Seed, B. (1987) Nature 329:840 or Kaufman et al. (1987) EMBO J. 6:187-195). Promoters preferably to be used in this context are of viral origin such as, for example, promoters of polyoma virus, adenovirus 2, cytomegalovirus or simian virus 40. Further prokaryotic and eukaryotic expression systems are mentioned 20 in Chapter 16 and 17 in Sambrook et al., Molecular Cloning: A Laboratory Manual. 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989. Further advantageous vectors are described in Hellens et al. (Trends in plant science, 5, 2000).

25 The transgenic organisms which comprise a NCLP sequence are claimed within the scope of the present invention.

All of the above-described embodiments of the transgenic organisms, which comprise at least one nucleic acid sequence according to the invention come under the term 30 "transgenic organism according to the invention".

The present invention furthermore relates to the use of CLP, in a method for identifying herbicidally active test compounds.

35 The method according to the invention for identifying herbicidally active compounds preferably comprises the following steps:

- 40 i. bringing CLP into contact with one or more test compounds under conditions which permit the test compound(s) to bind to a nucleic acid sequence according to the invention or to CLP, and
- ii. detecting whether the test compound binds to the CLP of i), or

- iii. detecting whether the test compound reduces or blocks the enzymatic or biological activity of CLP of i), or
- 5 iv. detecting whether the test compound reduces or blocks the transcription, translation or expression of CLP of i).

The detection in accordance with step (ii) of the above method can be effected using techniques which identify the interaction between the polypeptide and ligand. In this
10 context, either the test compound or the enzyme can contain a detectable label such as, for example, a fluorescent label, a radioisotope, a chemiluminescent label or an enzyme label. Examples of enzyme labels are horseradish peroxidase, alkaline phosphatase or luciferase. The subsequent detection depends on the label and is known to the skilled worker.

15

In this context, five preferred embodiments which are also suitable for high-throughput methods (HTS) in connection with the present invention must be mentioned in particular:

- 20 1. The average diffusion rate of a fluorescent molecule as a function of the mass can be determined in a small sample volume via fluorescence correlation spectroscopy (FCS) (Proc. Natl. Acad. Sci. USA (1994) 11753-11755). FCS can be employed for determining protein/ligand interactions by measuring the change in the mass, or the changed diffusion rate which this entails, of a test compound
25 when binding to CLP. A method according to the invention can be designed directly for measuring the binding of a test compound labeled by a fluorescent molecule. As an alternative, the method according to the invention can be designed in such a way that a chemical reference compound which is labeled by a fluorescent molecule is displaced by further test compounds ("displacement assay").
- 30 2. Fluorescence polarization exploits the characteristic of a quiescent fluorophore excited with polarized light to likewise emit polarized light. If, however, the fluorophore is allowed to rotate during the excited state, the polarization of the fluorescent light which is emitted is more or less lost. Under otherwise identical conditions (for example temperature, viscosity, solvent), the rotation is a function of molecule size, whereby findings regarding the size of the fluorophore-bound residue can be obtained via the reading (Methods in Enzymology 246 (1995), pp. 283-300). A method according to the invention can be designed directly for
35 measuring the binding of a test compound labeled with a fluorescent molecule to the CLP. As an alternative, the method according to the invention may also take the form of the "displacement assay" described under 1.
- 40

3. Fluorescence resonance energy transfer (FRET) is based on the irradiation-free energy transfer between two spatially adjacent fluorescent molecules under suitable conditions. A prerequisite is that the emission spectrum of the donor molecule overlaps with the excitation spectrum of the acceptor molecule. The fluorescent label of CLP, and binding test compound, the binding can be measured by means of FRET (Cytometry 34, 1998, pp. 159-179). As an alternative, the method according to the invention may also take the form of the "displacement assay" described under 1. An especially suitable embodiment of FRET technology is "Homogeneous Time Resolved Fluorescence" (HTRF) as can be obtained from Packard BioScience.
4. Surface-enhanced laser desorption/ionization (SELDI) in combination with a time-of-flight mass spectrometer (MALDI-TOF) makes possible the rapid analysis of molecules on a support and can be used for analyzing protein/ligand interactions (Worral et al., (1998) Anal. Biochem. 70:750-756). In a preferred embodiment, CLP, is immobilized on a suitable support and incubated with the test compound. After one or more suitable wash steps, the test compound molecules which are additionally bound to CLP, can be detected by means of the abovementioned methodology and test compounds which are bound to CLP, can thus be selected.
5. The measurement of surface plasmon resonance is based on the change in the refractive index at a surface when a test compound binds to a protein which is immobilized to said surface. Since the change in the refractive index is identical for virtually all proteins and polypeptides for a defined change in the mass concentration at the surface, this method can be applied to any protein in principle (Lindberg et al. Sensor Actuators 4 (1983) 299-304; Malmquist Nature 361 (1993) 186-187). The measurement can be carried out for example with the automatic analyzer based on surface plasmon resonance which is available from Biacore (Freiburg) at a throughput of, currently, up to 384 samples per day. A method according to the invention can be designed directly for measuring the binding of a test compound to CLP. As an alternative, the method according to the invention may also take the form of the "displacement assay" described under 1.

The compounds identified via the abovementioned methods 1 to 5 may be suitable as inhibitors. All of the substances identified via the abovementioned methods can subsequently be checked for their herbicidal action in another embodiment of the method according to the invention.

Furthermore, there exists the possibility of detecting further candidates for herbicidal active ingredients by molecular modeling via elucidation of the three-dimensional struc-

ture of CLP, by x-ray structure analysis. The preparation of protein crystals required for x-ray structure analysis, and the relevant measurements and subsequent evaluations of these measurements, the detection of a binding site in the protein, and the prediction of potential inhibitor structures are known to the skilled worker. In principle, an optimization of the compound identified by the abovementioned methods is also possible via molecular modeling.

A preferred embodiment of the method according to the invention, which is based on steps i) and ii), consists in selecting a test compound which reduces or blocks the activity of the CLP. Preferably, the activity of the CLP, incubated with the test compound is herein compared with the activity of a CLP, not incubated with a test compound.

A more preferred embodiment of the method based on steps i) and ii) consists in

- i. expressing CLP in a transgenic organism according to the invention or growing an organism which naturally contains a CLP,
- ii. bringing CLP, of step i) in the cell digest of the transgenic or nontransgenic organism, in partially purified or in homogeneously purified form, into contact with a test compound; and
- iii. selecting a compound which reduces or blocks the activity of the nuclear encoded Clp Protease. Preferably the activity of CLP incubated with the test compound is herein compared with the activity of a CLP, not incubated with a test compound.

The solution containing the CLP, can consist of the lysate of the original organism or of the transgenic organism which has been transformed with an expression cassette according to the invention. If necessary, the CLP, can be purified partially or fully via customary methods. A general overview over current protein purification techniques is described, for example, in Ausubel, F.M. et al., Current Protocols in Molecular Biology, Greene Publishing Assoc. and Wiley-Interscience (1994); ISBN 0-87969-309-6. In the case of recombinant preparation, the protein which has been fused with an affinity tag can be purified via affinity chromatography as is known to the skilled worker.

The CLP, which is required for in vitro methods can thus be isolated either by means of heterologous expression from a transgenic organism according to the invention or from an organism containing CLP, for example from an undesired plant, the term "undesired plant" being understood as meaning the species mentioned at the outset.

To identify herbicidal compounds, the CLP, is now incubated with a test compound. After a reaction time, the enzymatic activity of the CLP, incubated with the test com-

36

pound is determined in comparison with a CLP, not incubated with a test compound. If the CLP, is inhibited, a significant decrease in activity in comparison with the activity of the noninhibited polypeptide according to the invention is observed, the result being a reduction of at least 10%, advantageously at least 20%, preferably at least 30%, especially preferably by at least 50%, up to 100% reduction (blocking). Preferred is an inhibition of at least 50% at test compound concentrations of 10^{-4} M, preferably at 10^{-5} M, especially preferably of 10^{-6} M, based on enzyme concentration in the micromolar range.

10 The enzymatic activity of CLP, can be determined for example by an activity assay in which the increase of the product, the decrease of the substrate (or starting material) or the decrease or increase of the cofactor are determined, or by a combination of at least two of the abovementioned parameters, as a function of a defined period of time.

15 Examples of suitable substrates are, for example small peptides and modified small peptides like peptides coupled to a fluorogenic molecule such as aminomethylcoumarin and succinylated peptides.

20 If appropriate, derivatives of the abovementioned compounds which contain a detectable label such as, for example, a fluorescent label (e.g. fluorogenic substrates such as N-Suc-Leu-Tyr-(7-amino-4-methylcoumarine) (SLT-AMC), Suc-Ala-Ala-Ala-AMC, Suc-Leu-Leu-Val-Tyr-AMC, Suc-Ala-Ala-Phe-AMC, Suc-Ile-Ile-Trp-AMC, Suc-Ala-Phe-Lys-AMC), a radioisotope label or a chemiluminescent label, may also be used.

25 The amounts of substrates to be employed in the activity tests may range between 0.5 and 100 mM, based on 1-100 µg/ml enzyme.

[

The activity can be determined for example by tracking Proteolysis fluorimetrically when using fluorogenic Peptide substrates analogously to the method described by

30 Woo et al. 1989 The Journal of Biological Chemistry 264, pp.2088-2091, which is herein incorporated by reference.

The activity may also be determined in an ATP-dependent fashion in the presence of ClpA, ClpB or ClpC Protein as described in Halperin et al. 2001, Planta 213, pp. 614-

35 619. The preferred Substrate is then b-casein.

Furthermore the activity may be measured by HPLC and HPLC-MS methods detecting fragments of the peptides used as substrates.

40 Another preferred embodiment of the method according to the invention which is based on steps i) and iii) consists of the following steps:

37

- i. generating a transgenic organism according to the invention comprising a nucleic acid sequence according to the invention, wherein CLP is expressed recombinantly;
- 5 ii. applying a test compound to the transgenic organism of i) and to a nontransgenic organism of the same species;
- iii. determining the growth or the viability of the transgenic and the nontransgenic organisms after application of the test substance, and
- 10 iv. selecting test compounds which bring about a reduced growth or a limited viability of the nontransgenic organism in comparison with the growth of the transgenic organism.
- 15 In this context, the difference in growth in step iv) for the selection of a herbicidally active inhibitor amounts to at least 10%, by preference 20%, preferably 30%, especially preferably 40% and very especially preferably 50%.

20 The transgenic organism in this context is preferably a plant, an alga, a cyanobacterium, for example of the genus *Synechocystis* or a proteobacterium such as, for example, *Magnetococcus* sp. MC1, preferably plants which can be transformed by means of customary techniques, such as *Arabidopsis thaliana*, *Allium cepa*, *Ananas comosus*, *Arachis hypogaea*, *Asparagus officinalis*, *Beta vulgaris* spec. *altissima*, *Beta vulgaris* spec. *rapa*, *Brassica napus* var. *napus*, *Brassica napus* var. *napobrassica*, *Brassica* 25 *rapa* var. *silvestris*, *Camellia sinensis*, *Carthamus tinctorius*, *Carya illinoensis*, *Citrus limon*, *Citrus sinensis*, *Coffea arabica* (*Coffea canephora*, *Coffea liberica*), *Cucumis sativus*, *Cynodon dactylon*, *Daucus carota*, *Elaeis guineensis*, *Fragaria vesca*, *Glycine max*, *Gossypium hirsutum*, (*Gossypium arboreum*, *Gossypium herbaceum*, *Gossypium vitifolium*), *Helianthus annuus*, *Hevea brasiliensis*, *Hordeum vulgare*, *Humulus lupulus*, 30 *Ipomoea batatas*, *Juglans regia*, *Lens culinaris*, *Linum usitatissimum*, *Lycopersicon lycopersicum*, *Malus* spec., *Manihot esculenta*, *Medicago sativa*, *Musa* spec., *Nicotiana tabacum* (*N. rustica*), *Olea europaea*, *Oryza sativa*, *Phaseolus lunatus*, *Phaseolus vulgaris*, *Picea abies*, *Pinus* spec., *Pisum sativum*, *Prunus avium*, *Prunus persica*, *Pyrus communis*, *Ribes sylvestre*, *Ricinus communis*, *Saccharum officinarum*, *Secale cereale*, 35 *Solanum tuberosum*, *Sorghum bicolor* (s. *vulgare*), *Theobroma cacao*, *Trifolium pratense*, *Triticum aestivum*, *Triticum durum*, *Vicia faba*, *Vitis vinifera*, *Zea mays*, or cyanobacteria which can be transformed readily, such as *Synechocystis*, into which the sequence encoding a polypeptide according to the invention has been incorporated by transformation. These transgenic organisms thus show increased tolerance to compounds which inhibit the polypeptide according to the invention. "Knock-out" mutants in 40 which the analogous CLProtease gene which is naturally present in this organism has been selectively switched off may also be used.

However, the abovementioned embodiment of the method according to the invention can also be used for identifying substances with a growth-regulatory action. In this context, the transgenic organism employed is a plant. The method for identifying substances with growth-regulatory activity thus comprises the following steps:

- i. generating a transgenic plant comprising a nucleic acid sequence according to the invention encoding CLP, wherein CLP is expressed recombinantly;
- 10 ii. applying a test substance to the transgenic plant of i) and to a nontransgenic plant of the same variety,
- iii. determining the growth or the viability of the transgenic plant and the nontransgenic plant after application of the test compound, and
- 15 iv. selecting test substances which bring about a reduced growth of the nontransgenic plant in comparison with the growth of the transgenic plant.

Here, step iv) involves the selection of test compounds which bring about a modified growth of the nontransgenic organism in comparison with the growth of the transgenic organism. Modified growth is understood as meaning, in this context, inhibition of the vegetative growth of the plants, which can manifest itself in particular in reduced longitudinal growth. Accordingly, the treated plants show stunted growth; moreover, their leaves are darker in color. In addition, modified growth is also understood as meaning a change in the course of maturation over time, the inhibition or promotion of lateral branched growth of the plants, shortened or extended developmental stages, increased standing ability, the growth of larger amounts of buds, flowers, leaves, fruits, seed kernels, roots and tubers, an increased sugar content in plants such as sugarbeet, sugar cane and citrus fruit, an increased protein content in plants such as cereals or soybean, or stimulation of the latex flow in rubber trees. The skilled worker is familiar with the detection of such modified growth.

It is also possible, in the method according to the invention, to employ a plurality of test compounds in a method according to the invention. If a group of test compounds affect the target, then it is either possible directly to isolate the individual test compounds or to divide the group of test compounds into a variety of subgroups, for example when it consists of a multiplicity of different components, in order to thus reduce the number of the different test compounds in the method according to the invention. The method according to the invention is then repeated with the individual test compound or the relevant subgroup of test compounds. Depending on the complexity of the sample, the above-described steps can be carried out repeatedly, preferably until the subgroup

identified in accordance with the method according to the invention only comprises a small number of test compounds, or indeed just one test compound.

5 All of the above-described methods for identifying inhibitors with herbicidal or growth-regulatory activity are hereinbelow referred to as "methods according to the invention".

10 All of the compounds which have been identified via the methods according to the invention can subsequently be tested in vivo for their herbicidal and growth-regulatory activity. One possibility of testing the compounds for herbicidal action is to use duckweed, *Lemna minor*, in microtiter plates. Parameters which can be measured are changes in the chlorophyll content and the photosynthesis rate. It is also possible to apply the compound directly to undesired plants, it being possible to identify the herbicidal action for example via restricted growth.

15 The method according to the invention can advantageously also be carried out in high-throughput methods, known as HTS, which makes possible the simultaneous testing of a multiplicity of different compounds.

20 The use of supports which contain one or more of the nucleic acid molecules according to the invention, one or more of the vectors containing the nucleic acid sequence according to the invention, one or more transgenic organisms containing at least one of the nucleic acid sequences according to the invention or one or more (poly)peptides encoded via the nucleic acid sequences according to the invention lends itself to carrying out HTS in practice.

25 Supports which contain one or more of the NCLP sequences, one or more of the vectors comprising the NCLP sequences one or more transgenic organisms containing at least one NCLP sequences or one or more (poly)peptides encoded by the NCLP sequences are part of the present invention.

30 The support used can be solid or liquid, but is preferably solid and especially preferably a microtiter plate. The abovementioned supports also form part of the subject matter of the present invention. In accordance with the most widely used technique, 96-well, 384-well and 1536-well microtiter plates which, as a rule, can comprise volumes of 200 35 μ l, are used. Besides the microtiter plates, the further components of an HTS system which match the corresponding microtiter plates, such as a large number of instruments, materials, automatic pipetting devices, robots, automated plate readers and plate washers, are commercially available.

40 In addition to the HTS systems based on microtiter plates, what are known as "free-format assays" or assay systems where no physical barriers exist between the samples, as described, for example, in Jayaickreme et al., Proc. Natl. Acad. Sci U.S.A. 19

(1994) 161418; Chelsky, "Strategies for Screening Combinatorial Libraries", First Annual Conference of The Society for Biomolecular Screening in Philadelphia, Pa. (Nov. 7-10, 1995); Salmon et al., Molecular Diversity 2 (1996), 5763 and US 5,976,813, may also be used.

5

The invention furthermore relates to herbicidally active compounds identified by the methods according to the invention. These compounds are hereinbelow referred to as "selected compounds". They have a molecular weight of less than 1000 g/mol, advantageously less than 500 g/mol, preferably less than 400 g/mol, especially preferably less than 300 g/mol. Herbicidally active compounds have a K_i value of less than 1 mM, preferably less than 1 μ M, especially preferably less than 0.1 μ M, very especially preferably less than 0.01 μ M.

10

15 The invention furthermore relates to compounds with growth-regulatory activity identified by the methods according to the invention. These compounds too are hereinbelow referred to as "selected compounds".

15

Naturally, the selected compounds can also be present in the form of their agriculturally useful salts. Agriculturally useful salts which are suitable are mainly the salts of those cations, or the acid addition salts of those acids, whose cations, or anions, do not adversely affect the herbicidal action of the herbicidally active compounds identified via the methods according to the invention.

20

25 If the selected compounds contain asymmetrically substituted α -carbon atoms, they may furthermore also be present in the form of racemates, enantiomer mixtures, pure enantiomers or, if they have chiral substituents, also in the form of diastereomer mixtures.

25

The selected compounds can be chemically synthesized substances or substances produced by microbes and can be found, for example, in cell extracts of, for example, plants, animals or microorganisms. The reaction mixture can be a cell-free extract or comprise a cell or cell culture. Suitable methods are known to the skilled worker and are described generally for example in Alberts, Molecular Biology the cell, 3rd Edition (1994), for example chapter 17. The selected compounds may also originate from comprehensive substance libraries.

30

35

Candidate test compounds can be expression libraries such as, for example, cDNA expression libraries, peptides, proteins, nucleic acids, antibodies, small organic substances, hormones, PNAs or the like (Milner, *Nature Medicine* 1 (1995), 879–880; Hupp, *Cell*. 83 (1995), 237–245; Gibbs, *Cell*. 79 (1994), 193–198 and references cited therein).

40

The selected compounds can be used for controlling undesired vegetation and/or as growth regulators. Herbicidal compositions comprising the selected compounds afford very good control of vegetation on noncrop areas. In crops such as wheat, rice, maize, soybean and cotton, they act against broad-leaved weeds and grass weeds without
5 inflicting any significant damage on the crop plants. This effect is observed in particular at low application rates. The selected compounds can be used for controlling the harmful plants which have already been mentioned above.

Depending on the application method in question, selected compounds, or herbicidal
10 compositions comprising them, can advantageously also be employed in a further number of crop plants for eliminating undesired plants. Examples of suitable crops are:

Allium cepa, *Ananas comosus*, *Arachis hypogaea*, *Asparagus officinalis*, *Beta vulgaris* spec. *altissima*, *Beta vulgaris* spec. *rapa*, *Brassica napus* var. *napus*, *Brassica napus* var. *napobrassica*, *Brassica rapa* var. *silvestris*, *Camellia sinensis*, *Carthamus tinctorius*, *Carya illinoensis*, *Citrus limon*, *Citrus sinensis*, *Coffea arabica* (*Coffea canephora*, *Coffea liberica*), *Cucumis sativus*, *Cynodon dactylon*, *Daucus carota*, *Elaeis guineensis*, *Fragaria vesca*, *Glycine max*, *Gossypium hirsutum*, (*Gossypium arboreum*, *Gossypium herbaceum*, *Gossypium vitifolium*), *Helianthus annuus*, *Hevea brasiliensis*,
15 *Hordeum vulgare*, *Humulus lupulus*, *Ipomoea batatas*, *Juglans regia*, *Lens culinaris*, *Linum usitatissimum*, *Lycopersicon lycopersicum*, *Malus spec.*, *Manihot esculenta*, *Medicago sativa*, *Musa spec.*, *Nicotiana tabacum* (*N. rustica*), *Olea europaea*, *Oryza sativa*, *Phaseolus lunatus*, *Phaseolus vulgaris*, *Picea abies*, *Pinus spec.*, *Pisum sativum*, *Prunus avium*, *Prunus persica*, *Pyrus communis*, *Ribes sylvestre*, *Ricinus communis*, *Saccharum officinarum*, *Secale cereale*, *Solanum tuberosum*, *Sorghum bicolor* (s. *vulgare*), *Theobroma cacao*, *Trifolium pratense*, *Triticum aestivum*, *Triticum durum*,
20 *Vicia faba*, *Vitis vinifera*, *Zea mays*.

In addition, the selected compounds can also be used in crops which tolerate the action of herbicides owing to breeding, including recombinant methods. The generation of
30 such crops is described hereinbelow.

The invention furthermore relates to a method of preparing the herbicidal or growth-regulatory composition which has already been mentioned above, which comprises
35 formulating selected compounds with suitable auxiliaries to give crop protection products.

The selected compounds can be formulated for example in the form of directly sprayable aqueous solutions, powders, suspensions, also highly concentrated aqueous, oily
40 or other suspensions or suspoemulsions or dispersions, emulsifiable concentrates, emulsions, oil dispersions, pastes, dusts, materials for spreading or granules, and applied by means of spraying, atomizing, dusting, spreading or pouring. The use forms

depend on the intended use and the nature of the selected compounds; in any case, they should guarantee the finest possible distribution of the selected compounds. The herbicidal compositions comprise a herbicidally active amount of at least one selected compound and auxiliaries conventionally used in the formulation of herbicidal compositions.

For the preparation of emulsions, pastes or aqueous or oily formulations and dispersible concentrates (DC), the selected compounds can be dissolved or dispersed in an oil or solvent, it being possible to add further formulation auxiliaries for homogenization.

However, it is also possible to prepare liquid or solid concentrates from selected compound, if appropriate solvents or oil and, optionally, further auxiliaries comprising liquid or solid concentrates, and these concentrates are suitable for dilution with water. The following can be mentioned: emulsifiable concentrates (EC, EW), suspensions (SC), soluble concentrates (SL), dispersible concentrates (DC), pastes, pills, wettable powders or granules, it being possible for the solid formulations either to be soluble or dispersible (wetable) in water. In addition, suitable powders or granules or tablets can additionally be provided with a solid coating which prevents abrasion or premature release of the active ingredient.

In principle, the term "auxiliaries" is understood as meaning the following classes of compounds: antifoams, thickeners, wetting agents, tackifiers, dispersants, emulsifiers, bactericides and/or thixotropic agents. The skilled worker is familiar with the meaning of the abovementioned agents.

SLs, EWs and ECs can be prepared by simply mixing the ingredients in question; powders can be prepared by mixing or grinding in specific types of mills (for example hammer mills). DCs, SCs and SEs are usually prepared by wet milling, it being possible to prepare an SE from an SC by addition of an organic phase which may comprise further auxiliaries or selected compounds. The preparation is known. Powders, materials for spreading and dusts can advantageously be prepared by mixing or cogrinding the active substances together with a solid carrier. Granules, for example coated granules, impregnated granules and homogeneous granules, can be prepared by binding the selected compounds to solid carriers. The skilled worker is familiar with further details regarding their preparation, which are mentioned for example in the following publications: US 3,060,084, EP-A 707445 (for liquid concentrates), Browning, "Agglomeration", Chemical Engineering, Dec. 4, 1967, 147-48, Perry's Chemical Engineer's Handbook, 4th Ed., McGraw-Hill, New York, 1963, pages 8-57 and et seq. WO 91/13546, US 4,172,714, US 4,144,050, US 3,920,442, US 5,180,587, US 5,232,701, US 5,208,030, GB 2,095,558, US 3,299,566, Klingman, Weed Control as a Science, John Wiley and Sons, Inc., New York, 1961, Hance et al., Weed Control Handbook, 8th Ed., Blackwell Scientific Publications, Oxford, 1989 and Mollet, H., Grubemann, A., Formu-

lation technology, Wiley VCH Verlag GmbH, Weinheim (Federal Republic of Germany), 2001.

5 The skilled worker is familiar with a multiplicity of inert liquid and/or solid carriers which are suitable for the formulations according to the invention, such as, for example, liquid additives such as mineral oil fractions of medium to high boiling point such as kerosene or diesel oil, furthermore coal tar oils and oils of vegetable or animal origin, aliphatic, cyclic and aromatic hydrocarbons, for example paraffin, tetrahydrophthalene, alkylated naphthalenes or their derivatives, alkylated benzenes or their derivatives, alcohols such
10 as methanol, ethanol, propanol, butanol and cyclohexanol, ketones such as cyclohexanone, or strongly polar solvents, for example amines such as N-methylpyrrolidone or water.

15 Examples of solid carriers are mineral earths such as silicas, silica gels, silicates, talc, kaolin, limestone, lime, chalk, bole, loess, clay, dolomite, diatomaceous earth, calcium sulfate, magnesium sulfate, magnesium oxide, ground synthetic materials, fertilizers such as ammonium sulfate, ammonium phosphate, ammonium nitrate, ureas and products of vegetable origin such as cereal meal, tree bark meal, wood meal and nut-shell meal, cellulose powders or other solid carriers.

20 The skilled worker is familiar with the multiplicity of surface-active substances (surfactants) which are suitable for the formulations according to the invention such as, for example, alkali metal salts, alkaline earth metal salts or ammonium salts of aromatic sulfonic acids for example lignosulfonic acid, phenolsulfonic acid, naphthalenesulfonic
25 acid, and dibutylnaphthalenesulfonic acid, and of fatty acids, of alkyl- and alkylarylsulfonates, of alkyl sulfates, lauryl ether sulfates and fatty alcohol sulfates, and salts of sulfated hexa-, hepta- and octadecanols and of fatty alcohol glycol ethers, condensates of sulfonated naphthalene and its derivatives with formaldehyde, condensates of naphthalene or of the naphthalenesulfonic acids with phenol and formaldehyde, poly-
30 oxyethylene octylphenol ether, ethoxylated isooctyl-, octyl- or nonylphenol, alkylphenyl polyglycol ethers, tributylphenyl polyglycol ether, alkylaryl polyether alcohols, isotridecyl alcohol, fatty alcohol/ethylene oxide condensates, ethoxylated castor oil, polyoxyethylene alkyl ethers or polyoxypropylene alkyl ethers, lauryl alcohol polyglycol ether acetate, sorbitol esters, lignosulfite waste liquors or methylcellulose.

35 The herbicidal compositions, or the selected compounds, can be applied pre- or post-emergence. If the selected compounds are less well tolerated by certain crop plants, application techniques may be used in which the selected compounds are sprayed, with the aid of the spraying apparatus, in such a way that they come into as little contact, if any, with the leaves of the sensitive crop plants while the selected compounds reach the leaves of undesired plants which grow underneath, or the bare soil surface
40 (post-directed, lay-by).

Depending on the intended purpose of the control measures, the season, the target plants and the growth stage, the application rates of selected compounds amount to 0.001 to 3.0, preferably 0.01 to 1.0 kg/ha.

5

The invention is illustrated in greater detail by the examples which follow, which are not to be considered as limiting.

General DNA manipulation and cloning methods

10

Cloning methods such as, for example, restriction cleavages, agarose gel electrophoreses, purification of DNA fragments, transfer of nucleic acids to nitrocellulose and nylon membranes, linking DNA fragments, transformation of *Escherichia coli* cells, growing bacterium and sequence analyses of recombinant DNA were carried out as described by Sambrook et al. (1989) (Cold Spring Harbor Laboratory Press: ISBN 0-87969-309-6) and Ausubel, F.M. et al., Current Protocols in Molecular Biology, Greene Publishing Assoc. and Wiley-Interscience (1994); ISBN 0-87969-309-6.

15

Molecular-biological standard methods for plants and plant transformation methods are described in Schultz et al., Plant Molecular Biology Manual, Kluwer Academic Publishers (1998), Reither et al., Methods in Arabidopsis Research, World scientific press (1992) and Arabidopsis: A Laboratory Manual (2001), ISBN 0-87969-573-0.

20

The bacterial strains used hereinbelow (*E. coli* DH5, XL-1 blue) were obtained from Stratagene, BRL Gibco or Invitrogen, Carlsberg, CA. The vectors used for cloning were pUC 19 from Amersham Pharmacia (Freiburg) and the vector pBinAR (Höfgen and Willmitzer, Plant Science 66, 1990, 221-230).

25

Example 1: Generation of a cDNA library in the plant transformation vector

30

To generate a cDNA library (hereinbelow termed "binary cDNA library") in a vector which can be used directly for transforming plants, mRNA was isolated from a variety of plant tissues and transcribed into double-stranded cDNA using the cDNA Synthese Kit (Amersham Pharmacia Biotech, Freiburg). The cDNA first-strand synthesis was carried out using T12-18 oligonucleotides following the manufacturer's instructions. After size fractionation and the ligation of EcoRI-NotI adapters following the manufacturer's instructions and filling up the overhangs with Pfu DNA polymerase (Stratagene), the cDNA population was normalized. The method of Kohci et al, 1995, Plant Journal 8, 771-776 was followed, the cDNA being amplified by PCR with the oligonucleotide N1 under the conditions given in Table 1.

35

40

Table 1

Temperature [°C]	Time [sec]	Number of cycles
94	300	1
94	8	10
52	60	
72	180	
94	8	10
50	60	
72	180	
94	8	10
48	60	
72	180	
72	420	1

- 5 The resulting PCR product was bound to the column matrix of the PCR purification kit (Qiagen, Hilden) and eluted with 300 mM NaP buffer, pH 7.0, 0.5 mM EDTA, 0.04% SDS. The DNA was denatured for 5 minutes in a boiling water bath and subsequently renatured for 24 hours at 60°C. 50 µl of the DNA were applied to a hydroxylapatite column and the column was washed 3 times with 1 ml of 10 mM NaP buffer, pH 6.8. The bound single-stranded DNA was eluted with 130 mM NaP buffer, pH 6.8, precipitated with ethanol and dissolved in 40 µl of water. 20 µl of growth were used for a further PCR amplification as described above. After further ssDNA concentration, a third PCR amplification was carried out as described above.
- 10
- 15 The plant transformation vector for taking up the cDNA population which had been generated as described above was generated via restriction enzyme cleavage of the vector pUC18 with SbfI and BamHI, purification of the vector fragment followed by filling up the overhangs with Pfu DNA polymerase and relegation with T4 DNA ligase (Stratagene). The resulting construct is hereinbelow termed pUC18SbfI-.
- 20 The vector pBinAR was first cleaved with NotI, the ends were filled up and the vector was relegated, cleaved with SbfI, the ends were filled up and the vector was relegated and subsequently cleaved with EcoRI and HindIII. The resulting fragment was ligated into a derivative of the binary plant transformation vector pPZP (Hajdukiewicz, P, Svab, Z, Maliga, P., (1994) Plant Mol Biol 25:989-994) which makes possible the transformation of plants by means of agrobacterium and mediates kanamycin resistance in transgenic plants. The construct generated thus is hereinbelow termed pSun12/35S.
- 25

pUC18SbfI- was used as template in a polymerase chain reaction (PCR) with the oligonucleotides V1 and V2 (see Table 2) and Pfu DNA polymerase. The resulting frag-

ment was ligated into the SmaI-cut pSun12/35S, giving rise to pSunblues2. Following cleavage with NotI, dephosphorylation with shrimp alkaline phosphatase (Roche Diagnostics, Mannheim) and purification of the vector fragment, pSunblues2 was ligated with the normalized, likewise NotI-cut cDNA population. Following transformation into *E. coli* XI-1blue (Stratagene), the resulting clones were deposited into microtiter plates. The binary cDNA library contains cDNAs in "sense"- and in "antisense" orientation under the control of the cauliflower mosaic virus 35S promoter, and, after transformation into tobacco plants, these cDNAs can, accordingly, lead to "cosuppression" and "antisense" effects.

Table 2: Oligonucleotides used

Oligonucleotide	Nucleic acid sequence
N1	5'-AGAATTCGCGGCCGCT-3' (SEQ ID NO:23)
V1 (PWL93not)	5'-CTCATGCGGCCGCGCGCAACGCAATTAATGTG-3' (SEQ ID NO:24)
V2 (pWL92)	5'-TCATGCGGCCGCGAGATCCAGTTCGATGTAAC-3' (SEQ ID NO:25)
G1 (35S)	5'-GTGGATTGATGTGATATCTCC-3' (SEQ ID NO:26)
G2 (OCS)	5'-GTAAGGATCTGAGCTACACAT-3' (SEQ ID NO:27)

Example 2: Transformation and analysis of tobacco plants

Selected clones of the binary cDNA library were transformed into *Agrobacterium tumefaciens* C58C1:pGV2260 and (Deblaere et al., Nucl. Acids. Res. 13(1984), 4777-4788) and incubated with Streptomycin/Spectinomycin selection. The material used for the transformation of tobacco plants (*Nicotiana tabacum* cv. Samsun NN) with one of the binary clones as depicted in table 3 was an overnight culture of a positively transformed agrobacterial colony diluted with YEB medium to OD600 = 0.8-1.6. Leaf discs of sterile plants (approx. 1 cm² each) were incubated for 5-10 minutes with a 1:50 agrobacterial dilution in a Petri dish. This was followed by incubation in the dark for 2 days at 25°C on Murashige-Skoog medium (Physiol. Plant. 15(1962), 473) supplemented with 2% sucrose (2MS medium) and 0.8% Bacto agar. The cultivation was continued after 2 days at a 16-hour-light/8-hour-darkness photoperiod and continued in a weekly rhythm on MS medium supplemented with 500mg/l Claforan (cefotaxime sodium), 50mg/l kanamycin, 1mg/l benzylaminopurin (BAP), 0.2mg/l naphthylacetic acid and 1.6g/l glucose. Regenerated shoots were transferred onto an MS medium supplemented with kanamycin and Claforan. Transgenic plants of lines as depicted in table 3 were generated in this manner.

Table 3: Plant lines generated

Partial cDNA with phenotype in transgenic tobacco	Plant line	Corresponding full length cDNA	Function
SEQ ID NO: 20	E_0000013511	SEQ ID NO: 3	ClpP2 protease
SEQ ID NO:19	E_0000008893	SEQ ID NO:11	ClpP5=ClpR1 protease
SEQ ID NO:21	E_0000012393	-	ClpP6 protease
-	-	SEQ ID NO:17	ClpP-like protease

The integration of the clone cDNA into the genome of the transgenic lines was detected via PCR with the oligonucleotides G1 and G2 (see Table 2) and genomic DNA prepared from the transgenic lines in question. To this end, TAKARA Taq DNA polymerase was preferably employed for this purpose, following the manufacturer's instructions (MoBiTec, Göttingen). The cDNA clone of the binary cDNA library, which clone had been used for the transformation, acted as template for a PCR reaction as the positive control. PCR products with an identical size or, if appropriate, identical cleavage patterns which were obtained after cleavage with a variety of restriction enzymes acted as proof that the corresponding cDNA had been integrated. In this manner, the insert of clones were detected in the respective transgenic plant lines (as depicted in table 3) with the belowmentioned phenotypes.

After the shoots had been transferred into soil, the plants were observed for 2-20 weeks in the greenhouse for the manifestation of phenotypes. It emerged that transgenic plants of lines E_0000012393, E_0000013511 and E_0000008893 were similar in phenotype. The plants showed severe chlorosis and concomitant growth retardation with respect to wild type plants after 2 weeks.

Example 3: Sequence analysis of the clones

SEQ ID NO:19 was fully sequenced and used for the detection of the corresponding full length clone SEQ ID NO:11. SEQ ID NO:11 is identical to nt002050074r-SEQ ID NO:19 in the overlapping region. An open reading frame of 867 nt (pos. 2-1162) encodes for 387 amino acids (SEQ ID NO: 4) with highest identity to ClpR1 from *Arabidopsis thaliana*. Sequence homology suggests that the 5'-ends of SEQ ID NO:11 and ClpR1 from *Arabidopsis thaliana* are very diverse and that nt006066004r is close to being full size with respect to ClpR1. MS-Analysis of isolated ClpP Proteins from *Arabidopsis* indicate, that the mature ClpR1 is several kDa shorter as suggested by the cDNA Sequence (Peltier et al. 2001, The Journal of Biological Chemistry 276, 99.16318-16327).

SEQ ID NO:20 was fully sequenced and used for the detection of the corresponding full length clone, SEQ ID NO: 3. SEQ ID NO:3 is identical to SEQ ID NO:20 in the overlap-

ping region. An open reading frame of 867 nt (pos. 11-877) encodes for 289 amino acids (SEQ ID NO:4) with highest identity to ClpP2 from *Arabidopsis thaliana*.

5 SEQ ID NO:21 was fully sequenced. The partial cDNA Sequence of 602 nt contains an open reading frame of 186 nt (nt 8-193) encoding for 62 amino acids (SEQ ID NO:22). This partial polypeptide shows highest identity to ClpP6 from *Arabidopsis thaliana* (SEQ ID NO:9)

10 A further ClpP-homolog cDNA of 906 nt (SEQ ID NO:17) was identified. An open reading frame of 711 nt (pos. 45-755) encodes for 237 amino acids (SEQ ID NO:18) with highest identity to a ClpP-like protein from *Arabidopsis thaliana* (GeneBank Acc. No. AK118523).

15 Thus, it was shown for the first time and in a surprising manner that the natural expression of nuclear encoded Clp protease encoding genes is essential for plants and that reduced expression leads to damage as depicted by the phenotypes mentioned in Example 2 demonstrating the suitability of nuclear encoded Clp-proteases as target for herbicides.

20 Example 4: Expression in *E.coli*

In order to generate active protein with nuclear encoded Clp-protease activity fragments of SEQ ID NO:11, -SEQ ID NO:3 and of SEQ ID NO:9 were subcloned into the expression vector pQE60 (Quiagen, Hilden, Germany). To this end the oligonucleotides
25 displayed in tab. 4 were used to amplify via polymerase chain reaction cDNA fragments that contain NcoI and BglII restriction sites. The PCR was carried out in 36 cycles following standard conditions (for example as described by Sambrook, J. et al. (1989) "Molecular cloning: A laboratory manual", Cold Spring Harbor Laboratory Press), the annealing temperatures being between 45 and 55°C and the polymerization
30 time being in each case 60 seconds per 1000bp. Cutting the cDNA fragments with NcoI and BglII restriction enzymes and ligation into pQE60 cut with the same enzymes delivered expression plasmids that were transformed into *E. coli*. Expression was performed in *E. coli* TOP 10F strains (Invitrogen, Karlsruhe, Germany) following induction with IPTG. Standard protocols (Invitrogen) were followed.

35 Enzyme preparations were achieved by breaking cells in a French-Press in 100mM Tris/HCl, pH 7.4, 2.5 mM EDTA, 1% Triton X-100.

40 The expression products were purified by affinity chromatography on Ni-agarose where appropriate. The manufacturer's instructions were followed (Qiagen).

Table 4

Construct	Primer (Nucleic acid sequence)
NT_ClpR1 ¹⁾	5'-TATACCATGGATTTGCCATCTTTG-3' (SEQ ID NO:28) 5'-ATAGATCTCACCTGGAGCCAG-3' (SEQ ID NO:29)
Nt_ClpP2 ¹⁾	5'-GAGCCCATGGCAAGAGGAG -3' (SEQ ID NO:30) 5'-ATAGATCTTTCTAGCTTGAACC-3' (SEQ ID NO:31)
AT_ClpP6 ²⁾	5'-TCAGCCATGGCCCTGGAGGAC -3' (SEQ ID NO:32) 5'-TAAGATCTTCAGTATTCTGTTTCC-3' (SEQ ID NO:33)

- 1) Template: *Nicotiana tabacum* cDNA library
- 2) Template: *Arabidopsis thaliana* cDNA library

5 Example 5: Activity assay

Isolated ClpP activity can be measured as described (Woo et al. 1989 The Journal of Biological Chemistry 264, pp.2088-2091) by using fluoregenic substrates such as N-Suc-Leu-Tyr-(7-amino-4-methylcoumarine) (SLT-AMC). The proteolytic cleavage delivers 7-amino-4-methylcoumarin, which can be detected fluorimetrically (emission at 460nm by excitation at 390 nm).

Standard assays contain: 50mM Tris/HCl, pH 8.0, 25mM MgCl₂, 1mM SLT-AMC and 1-100 µg ClpP Enzyme.

15 The assay is suitable in for high throughput screening in 96well and 384 well format.

Sequence Listing

20	Sequence	Function	Organism
	SEQ ID NO:1 (nucleic acid sequence)	ClpP1	Arabidopsis thaliana
	SEQ ID NO:2 (amino acid sequence)	ClpP1	Arabidopsis thaliana
25	SEQ ID NO:3 (nucleic acid sequence)	ClpP2	Nicotiana tabacum
	SEQ ID NO:4 (amino acid sequence)	ClpP2	Nicotiana tabacum
30	SEQ ID NO:5 (nucleic acid sequence)	ClpP3	Arabidopsis thaliana
	SEQ ID NO:6 (amino acid sequence)	ClpP3	Arabidopsis thaliana
35	SEQ ID NO:7 (nucleic acid sequence)	ClpP4	Arabidopsis thaliana

50

	SEQ ID NO:8 (amino acid sequence)	ClpP4	Arabidopsis thaliana
	SEQ ID NO:9 (nucleic acid sequence)	ClpP6	Arabidopsis thaliana
5	SEQ ID NO:10 (amino acid sequence)	ClpP6	Arabidopsis thaliana
	SEQ ID NO:11 (nucleic acid sequence)	ClpR1	Nicotiana tabacum
10	SEQ ID NO:12 (amino acid sequence)	ClpR1	Nicotiana tabacum
	SEQ ID NO:13 (nucleic acid sequence)	ClpR3	Arabidopsis thaliana
	SEQ ID NO:14 (amino acid sequence)	ClpR3	Arabidopsis thaliana
15	SEQ ID NO:15 (nucleic acid sequence)	ClpR4	Arabidopsis thaliana
	SEQ ID NO:16 (amino acid sequence)	ClpR4	Arabidopsis thaliana
20	SEQ ID NO:17 (nucleic acid sequence)	ClpP like	Arabidopsis thaliana
	SEQ ID NO:18 (amino acid sequence)	ClpP like	Arabidopsis thaliana
	SEQ ID NO:19 (nucleic acid sequence) (fragment)	ClpR1	Nicotiana tabacum
25	SEQ ID NO:20 (nucleic acid sequence) (fragment)	ClpP2	Nicotiana tabacum
30	SEQ ID NO:21 (nucleic acid sequence) (fragment)	ClpP6	Nicotiana tabacum
	SEQ ID NO:22 (amino acid sequence) (fragment)	ClpP6	Nicotiana tabacum
35	SEQ ID NO:23-33: Primer (nucleic acid sequences)		

We claim:

1. The use of nuclear encoded Clp-protease in a method for identifying herbicides.
2. The use as claimed in claim 1, wherein the Clp-protease is
 - a) selected from the group consisting of ClpP1-protease, ClpP2-protease, ClpP3-protease, ClpP4-protease and ClpP6-protease; or
 - b) selected from the group consisting of ClpR1-protease, ClpR3-protease, ClpR4-protease; or
 - c) ClpP-like-protease.
3. A plant nucleic acid sequence encoding a ClpP2-protease comprising:
 - a) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:3, or
 - b) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:4 by backtranslating, or
 - c) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:3 which has an identity with SEQ ID NO:3 of at least 66%.
4. A plant nucleic acid sequence encoding a ClpR1-protease comprising:
 - a) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:11, or
 - b) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:12 by backtranslating, or
 - c) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:11 which has an identity with SEQ ID NO:11 of at least 69%.
5. A plant nucleic acid sequence encoding a ClpP-like-protease comprising:

52

- a) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:17, or
- 5 b) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:18 by backtranslating, or
- c) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:17 which has an identity with SEQ ID NO:17 of has at least 67%.
- 10 6. A polypeptide with the activity of a nuclear encoded Clp-protease, encoded by a nucleic acid molecule as claimed in claim 3, 4 or 5.
- 15 7. An expression cassette comprising
- a) genetic control sequences in operable linkage with a nucleic acid sequence as claimed in claim 3, 4 or 5; or
- 20 b) additional functional elements, or
- c) a combination of a) and b).
8. A vector comprising an expression cassette as claimed in claim 7.
- 25 9. A transgenic organism comprising at least one nucleic acid sequence as claimed in claim 4, 5 or 6 encoding a polypeptide with the activity of a Clp-protease, an expression cassette as claimed in claim 7 or a vector as claimed in claim 8, selected from among bacteria, yeasts, fungi, animal cells or plant cells.
- 30 10. A method for identifying substances with herbicidal activity, comprising the following steps:
- 35 i. bringing a nuclear encoded Clp-protease into contact with one or more test compounds under conditions which permit the test compound(s) to bind to the nucleic acid molecule encoding Clp-protease or to the nuclear encoded Clp-protease, and
- 40 ii. detecting whether the test compound binds to the Clp-protease of i), or
- iii. detecting whether the test compound reduces or blocks the enzymatic or biological activity of the Clp-protease of i), or

- iv. detecting whether the test compound reduces or blocks the transcription, translation or expression of the Clp-protease of i).

11. A method as claimed in claim 10, wherein the Clp-protease is

- a) selected from the group consisting of ClpP1-protease, ClpP2-protease, ClpP3-protease, ClpP4-protease and ClpP6-protease; or
- b) selected from the group consisting of ClpR1-protease, ClpR3-protease, ClpR4-protease; or
- c) ClpP-like-protease.

12. A method as claimed in claim 10, wherein

- a) the ClpP1-protease is encoded by a nucleic acid sequence which comprises:
- i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:1, or
- ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:2 by back translating, or
- iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:1 which has an identity with SEQ ID NO:1 of has at least 50%;
- b) the ClpP2-protease is encoded by a nucleic acid sequence which comprises:
- i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:3, or
- ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:4 by back translating, or
- iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:3 which has an identity with SEQ ID NO:3 of has at least 50%;

- 5 c) the ClpP3-protease is encoded by a nucleic acid sequence which comprises:
- 5 i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:5, or
- 10 ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:6 by back translating, or
- 15 iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:5 which has an identity with SEQ ID NO:5 of has at least 50%;
- 15 d) the ClpP4-protease is encoded by a nucleic acid sequence which comprises:
- 20 i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:7, or
- 25 ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:8 by back translating, or
- 30 iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:7 which has an identity with SEQ ID NO:7 of has at least 50%;
- 35 e) the ClpP6-protease is encoded by a nucleic acid sequence which comprises:
- 40 i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:9, or
- 45 ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:10 by back translating, or
- 50 iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:9 which has an identity with SEQ ID NO:9 of has at least 50%;
- 55 f) the ClpR1-protease is encoded by a nucleic acid sequence which comprises:

55

- 5
- i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:11, or
- ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:12 by back translating, or
- 10
- iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:11 which has an identity with SEQ ID NO:11 of has at least 50%;
- g) the ClpR3-protease is encoded by a nucleic acid sequence which comprises:
- 15
- i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:13, or
- ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:14 by back translating, or
- 20
- iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:13 which has an identity with SEQ ID NO:13 of has at least 50%;
- h) the ClpR4-protease is encoded by a nucleic acid sequence which comprises:
- 25
- i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:15, or
- 30
- ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:16 by back translating, or
- 35
- iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:15 which has an identity with SEQ ID NO:15 of has at least 50%;
- i) the ClpP like-protease is encoded by a nucleic acid sequence which comprises:
- 40
- i) a nucleic acid sequence with the nucleic acid sequence shown in SEQ ID NO:17, or

56

- ii) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO:18 by back translating, or
- 5 iii) a functional equivalent of nucleic acid sequence shown in SEQ ID NO:17 which has an identity with SEQ ID NO:17 of has at least 50%;
13. A method as claimed in claim 10, 11 or 12, wherein a test compound is selected which reduces or blocks the enzymatic or biological activity of Clp-protease.
- 10 14. A method as claimed in any of claims 10, 11, 12 or 13, wherein
- 15 i. either Clp-protease is expressed in a transgenic organism or an organism which naturally contains Clp-protease is grown,
- ii. the Clp-protease of step i) is brought into contact with a test compound in the cell digest of the transgenic or nontransgenic organism, in partially purified form or in homogeneously purified form, and
- 20 iii. selecting a test compound which reduces or blocks the enzymatic activity of the Clp-protease of step a).
15. A method as claimed in any of claims 10, 11, 12 or 13, which comprises the following steps:
- 25 i. generating a transgenic organism comprising a nucleic acid sequence encoding Clp-protease, wherein Clp-protease is expressed recombinantly;
- 30 ii. applying a test substance to the transgenic organism of i) and to a non-transgenic organism of the same genotype,
- iii. determining the growth or the viability of the transgenic plant and the non-transgenic plant after application of the test compound, and
- 35 iv. selecting test substances which bring about a reduced growth of the nontransgenic plant in comparison with the growth of the transgenic plant.
16. A method as claimed in claim 15, which is carried out in a plant organism, a cyanobacterium or proteobacterium.
- 40 17. A method for identifying substances with growth-regulatory activity, which comprises the following steps:

- i. generating a transgenic plant comprising a nucleic acid sequence Clp-protease, wherein Clp-protease is expressed recombinantly;
- 5 ii. applying a test substance to the transgenic plant of i) and to a nontransgenic plant of the same variety,
- iii. determining the growth or the viability of the transgenic plant and the non-transgenic plant after application of the test compound, and
- 10 iv. selecting test substances which bring about a reduced growth of the non-transgenic plant in comparison with the growth of the transgenic plant.
18. A method as claimed in any of claims 10 to 17, wherein the substances are identified in high-throughput screening method.
- 15
19. A support comprising one or more of the nucleic acid molecules as claimed in claim 3, 4, or 5 one or more expression cassettes as claimed in claim 7, one or more vectors as claimed in claim 8, one or more organisms as claimed in claim 9 or one or more (poly)peptides as claimed in claim 6.
- 20
20. A method as claimed in any of claims 10 to 18, wherein the substances are identified in High-Throughput Screening using a support as claimed in claim 19.
- 25
21. A compound with herbicidal activity, identified by one of the methods as claimed in any of claims 10 to 16, 18 and 20.
22. A compound with growth-regulatory activity, identified by the method as claimed in any of claims 17, 18 or 20.
- 30
23. A method for the preparation of an agrochemical composition, which comprises
- a) identifying a compound with herbicidal activity by one of the methods as claimed in any of claims 10 to 16, 18 and 20 or a compound with growth-regulatory activity as claimed in any of claims 17, 18 or 20, and
- 35
- b) formulating this compound together with suitable auxiliaries to give crop protection products with herbicidal or growth-regulatory activity.
- 40
24. A method for controlling undesired vegetation and/or for regulating the growth of plants, which comprises allowing at least one Clp-protease inhibitor as claimed in claim 21 or 22 or an agrochemical composition comprising a Clp-protease inhibi-

58

tor as claimed in claim 21 or 22 to act on plants, their environment and/or on seeds.

- 5 25. The use of a Clp-protease inhibitor as claimed in claim 21 or 22 or of an agro-chemical composition comprising a Clp-protease inhibitor as claimed in claim 21 or 22 in a method as claimed in claim 24.

Clp-protease as target for herbicides

5 The present invention relates to Clp-protease, which, when absent, brings about reduced growth and chlorotic leaves as target for herbicides. For this purpose, novel nucleic acid sequences encompassing SEQ ID NO:3, SEQ ID NO:11 and SEQ ID NO: 17 and functional equivalents of SEQ ID NO:3, SEQ ID NO:11 and SEQ ID NO: 17 are provided. Moreover, the present invention relates to the use of Clp-protease in a method for identifying compounds with herbicidal or growth-regulatory activity, and to the use of the compounds identified by this method as herbicides or growth regulators.

10

SEQUENCE LISTING

<110> BASF Aktiengesellschaft

5 <120> Clp-protease as target for herbicides

<130> 20030949

10 <160> 33

<170> PatentIn version 3.1

15 <210> 1
<211> 591
<212> DNA
<213> Arabidopsis thaliana

20 <220>
<221> CDS
<222> (1) .. (591)
<223>

25 <400> 1
atg cct att ggc gtt cca aaa gta cct ttt cga agt cct gga gaa gga 48
Met Pro Ile Gly Val Pro Lys Val Pro Phe Arg Ser Pro Gly Glu Gly
1 5 10 15

30 gat aca tct tgg gtt gac ata tac aac cga ctt tat cga gaa aga tta 96
Asp Thr Ser Trp Val Asp Ile Tyr Asn Arg Leu Tyr Arg Glu Arg Leu
20 25 30

35 ttt ttt tta ggc caa gag gtt gat acc gaa atc tog aat caa ctt att 144
Phe Phe Leu Gly Gln Glu Val Asp Thr Glu Ile Ser Asn Gln Leu Ile
35 40 45

40 agt ctt atg ata tat ctc agt ata gaa aag gat acc aaa gat ctt tat 192
Ser Leu Met Ile Tyr Leu Ser Ile Glu Lys Asp Thr Lys Asp Leu Tyr
50 55 60

45 ttg ttt ata aac tct cct ggt gga tgg gta ata tct gga atg gct att 240
Leu Phe Ile Asn Ser Pro Gly Gly Trp Val Ile Ser Gly Met Ala Ile
65 70 75 80

50 tat gat act atg caa ttt gtg cga ccc gat gta cag aca ata tgc atg 288
Tyr Asp Thr Met Gln Phe Val Arg Pro Asp Val Gln Thr Ile Cys Met
85 90 95

55 gga ttg gcc gct tca ata gca tcc ttt atc cta gtc gga gga gca att 336
Gly Leu Ala Ala Ser Ile Ala Ser Phe Ile Leu Val Gly Gly Ala Ile
100 105 110

60 acc aaa cgt ata gca ttc cct cac gct agg gta atg atc cat caa ccc 384
Thr Lys Arg Ile Ala Phe Pro His Ala Arg Val Met Ile His Gln Pro
115 120 125

65 gct agt tgc ttt tat gag gca caa acg gga gaa ttt atc ttg gaa gcg 432
Ala Ser Ser Phe Tyr Glu Ala Gln Thr Gly Glu Phe Ile Leu Glu Ala
130 135 140

gaa gaa tta ctt aaa ctt cgc gaa acc atc aca agg gtt tat gta caa 480
Glu Glu Leu Leu Lys Leu Arg Glu Thr Ile Thr Arg Val Tyr Val Gln
145 150 155 160

aga acg ggc aaa cct ata tgg gtt ata tcc gaa gac atg gaa cgg gat 528
Arg Thr Gly Lys Pro Ile Trp Val Ile Ser Glu Asp Met Glu Arg Asp

2

165 170 175

gtt ttt atg tca gca aca gaa gcc caa gct cat gga att gtt gat ctt 576
Val Phe Met Ser Ala Thr Glu Ala Gln Ala His Gly Ile Val Asp Leu
180 185 190

5

gta gag gtt caa taa 591
Val Ala Val Gln
195

10

<210> 2
<211> 196
<212> PRT
15 <213> Arabidopsis thaliana
<400> 2

20 Met Pro Ile Gly Val Pro Lys Val Pro Phe Arg Ser Pro Gly Glu Gly
1 5 10 15

25 Asp Thr Ser Trp Val Asp Ile Tyr Asn Arg Leu Tyr Arg Glu Arg Leu
20 25 30

30 Phe Phe Leu Gly Gln Glu Val Asp Thr Glu Ile Ser Asn Gln Leu Ile
35 40 45

35 Ser Leu Met Ile Tyr Leu Ser Ile Glu Lys Asp Thr Lys Asp Leu Tyr
50 55 60

40 Leu Phe Ile Asn Ser Pro Gly Gly Trp Val Ile Ser Gly Met Ala Ile
65 70 75 80

45 Tyr Asp Thr Met Gln Phe Val Arg Pro Asp Val Gln Thr Ile Cys Met
85 90 95

50 Gly Leu Ala Ala Ser Ile Ala Ser Phe Ile Leu Val Gly Gly Ala Ile
100 105 110

55 Thr Lys Arg Ile Ala Phe Pro His Ala Arg Val Met Ile His Gln Pro
115 120 125

60 Ala Ser Ser Phe Tyr Glu Ala Gln Thr Gly Glu Phe Ile Leu Glu Ala
130 135 140

65 Glu Glu Leu Leu Lys Leu Arg Glu Thr Ile Thr Arg Val Tyr Val Gln
145 150 155 160

Arg Thr Gly Lys Pro Ile Trp Val Ile Ser Glu Asp Met Glu Arg Asp
165 170 175

Val Phe Met Ser Ala Thr Glu Ala Gln Ala His Gly Ile Val Asp Leu
180 185 190

	Val Ala Val Gln	
	195	
5	<210> 3	
	<211> 1024	
	<212> DNA	
10	<213> Nicotiana tabacum	
	<220>	
	<221> CDS	
15	<222> (11)..(877)	
	<223>	
	<400> 3	
20	gcgggocgcta atg gog gtc act ttt ccg acc acc tct tcc tcg tat cta Met Ala Val Thr Phe Pro Thr Thr Ser Ser Ser Tyr Leu	49
	1 5 10	
25	cac tcg aga act aaa gtc cct cag cct tct tta agc tgc gcc agc aaa His Ser Arg Thr Lys Val Pro Gln Pro Ser Leu Ser Cys Ala Ser Lys	97
	15 20 25	
30	gtt ttt gtc gga tta aga agc caa tot cct aat tct tat ggg att gca Val Phe Val Gly Leu Arg Ser Gln Ser Pro Asn Ser Tyr Gly Ile Ala	145
	30 35 40 45	
35	gcg tct aat gta aat gtt gaa ttt cac aat aga gtg tac aga agt att Ala Ser Asn Val Asn Val Glu Phe His Asn Arg Val Tyr Arg Ser Ile	193
	50 55 60	
40	gaa tcc gga act aga gac agt aaa cca aca cgt gta cga gtt tcc atg Glu Ser Gly Thr Arg Asp Ser Lys Pro Thr Arg Val Arg Val Ser Met	241
	65 70 75	
45	atg ccc att ggg aca cca aga gta ccc tac aga aat oca act gag gga Met Pro Ile Gly Thr Pro Arg Val Pro Tyr Arg Asn Pro Thr Glu Gly	289
	80 85 90	
50	aca tgg cag tgg gtt gat ttg tgg aat got ctt tac cgt gaa cgt gtt Thr Trp Gln Trp Val Asp Leu Trp Asn Ala Leu Tyr Arg Glu Arg Val	337
	95 100 105	
55	att ttc atc gga caa cac ata gat gaa gaa ttt agc aac cag ata ttg Ile Phe Ile Gly Gln His Ile Asp Glu Glu Phe Ser Asn Gln Ile Leu	385
	110 115 120 125	
60	gca aca atg ctg tat ctt gac agt att gat gat tcc aag aag ctc tac Ala Thr Met Leu Tyr Leu Asp Ser Ile Asp Asp Ser Lys Lys Leu Tyr	433
	130 135 140	
65	ctg tat atc aat ggc cct ggt ggt gat cta act cca agc atg gcc atc Leu Tyr Ile Asn Gly Pro Gly Gly Asp Leu Thr Pro Ser Met Ala Ile	481
	145 150 155	
70	tac gac aca atg caa agt ctt aaa agt got gtt ggc acc cat tgt gtg Tyr Asp Thr Met Gln Ser Leu Lys Ser Ser Ala Val Gly Thr His Cys Val	529
	160 165 170	
75	ggc tat gcc tac aat ctt gcc ggt ttt ctt ctt got got gga gaa aag Gly Tyr Ala Tyr Asn Leu Ala Gly Phe Leu Leu Ala Ala Gly Glu Lys	577

4

	175	180	185	
5	ggc aat cga ttt gca atg cct ctt toa agg att gca cta caa tot cca Gly Asn Arg Phe Ala Met Pro Leu Ser Arg Ile Ala Leu Gln Ser Pro 190 195 200 205			625
10	gct gga gct ggc cgc gga cag gct gat gat att cgc aat gaa gca gat Ala Gly Ala Ala Arg Gly Gln Ala Asp Asp Ile Arg Asn Glu Ala Asp 210 215 220			673
	gaa ctt ctc aga att aga gat tac ctt ttc aag gag ttg gct gag aag Glu Leu Leu Arg Ile Arg Asp Tyr Leu Phe Lys Glu Leu Ala Glu Lys 225 230 235			721
15	aca ggc cag cct gtt gaa aag gtt cac aag gat tta agt cgg atg aag Thr Gly Gln Pro Val Glu Lys Val His Lys Asp Leu Ser Arg Met Lys 240 245 250			769
20	cga ctc aat gct caa gaa gct ctt gaa tat ggt ctt ata gac cgt ata Arg Leu Asn Ala Gln Glu Ala Leu Glu Tyr Gly Leu Ile Asp Arg Ile 255 260 265			817
25	gtt agg cct ccc cgt att aag gca gat gct cca cga aag gat acc aca Val Arg Pro Pro Arg Ile Lys Ala Asp Ala Pro Arg Lys Asp Thr Thr 270 275 280 285			865
	gca ggt ctt ggt tagtccatac acatcgtata atttatggct gatagtgggt Ala Gly Leu Gly			917
30	gtacgacttg cagtgttatt ttgcaatttc ttttgtttca totacatatt gaaotctttt gatctactta ttcaaaaaca tgaaatcctg agcagactag cggcgcg			977 1024
35	<210> 4 <211> 289 <212> PRT <213> Nicotiana tabacum			
40	<400> 4			
45	Met Ala Val Thr Phe Pro Thr Thr Ser Ser Ser Tyr Leu His Ser Arg 1 5 10 15			
	Thr Lys Val Pro Gln Pro Ser Leu Ser Cys Ala Ser Lys Val Phe Val 20 25 30			
50	Gly Leu Arg Ser Gln Ser Pro Asn Ser Tyr Gly Ile Ala Ala Ser Asn 35 40 45			
55	Val Asn Val Glu Phe His Asn Arg Val Tyr Arg Ser Ile Glu Ser Gly 50 55 60			
60	Thr Arg Asp Ser Lys Pro Thr Arg Val Arg Val Ser Met Met Pro Ile 65 70 75 80			
	Gly Thr Pro Arg Val Pro Tyr Arg Asn Pro Thr Glu Gly Thr Trp Gln 85 90 95			

5 Trp Val Asp Leu Trp Asn Ala Leu Tyr Arg Glu Arg Val Ile Phe Ile
100 105 110

10 Gly Gln His Ile Asp Glu Glu Phe Ser Asn Gln Ile Leu Ala Thr Met
115 120 125

15 Leu Tyr Leu Asp Ser Ile Asp Asp Ser Lys Lys Leu Tyr Leu Tyr Ile
130 135 140

20 Asn Gly Pro Gly Gly Asp Leu Thr Pro Ser Met Ala Ile Tyr Asp Thr
145 150 155 160

25 Met Gln Ser Leu Lys Ser Ala Val Gly Thr His Cys Val Gly Tyr Ala
165 170 175

30 Tyr Asn Leu Ala Gly Phe Leu Leu Ala Ala Gly Glu Lys Gly Asn Arg
180 185 190

35 Phe Ala Met Pro Leu Ser Arg Ile Ala Leu Gln Ser Pro Ala Gly Ala
195 200 205

40 Ala Arg Gly Gln Ala Asp Asp Ile Arg Asn Glu Ala Asp Glu Leu Leu
210 215 220

45 Arg Ile Arg Asp Tyr Leu Phe Lys Glu Leu Ala Glu Lys Thr Gly Gln
225 230 235 240

50 Pro Val Glu Lys Val His Lys Asp Leu Ser Arg Met Lys Arg Leu Asn
245 250 255

55 Ala Gln Glu Ala Leu Glu Tyr Gly Leu Ile Asp Arg Ile Val Arg Pro
260 265 270

60 Pro Arg Ile Lys Ala Asp Ala Pro Arg Lys Asp Thr Thr Ala Gly Leu
275 280 285

50 Gly

55 <210> 5
<211> 1124
<212> DNA
<213> Arabidopsis thaliana

60 <220>
<221> CDS
<222> (2)..(931)
<223>

<400> 5

5

10

15

20

25

30

35

10

50

EE

ttg aat cca tgg gag gcg aaa gaa tat ggt ttg atc gac gct gta atc 769
Leu Asn Pro Trp Glu Ala Lys Glu Tyr Gly Leu Ile Asp Ala Val Ile

7

	245	250	255	
5	gat gat ggg aaa cgg gga cta atc gct cca att gga gat ggt act cct Asp Asp Gly Lys Pro Gly Leu Ile Ala Pro Ile Gly Asp Gly Thr Pro 260 265 270			817
10	cct cct aaa acc aaa gtc tgg gat ctt tgg aaa gtc gaa gga acc aag Pro Pro Lys Thr Lys Val Trp Asp Leu Trp Lys Val Glu Gly Thr Lys 275 280 285			865
15	aaa gac aac act aac ttg cca tct gag cgc tcc atg aca cag aat ggt Lys Asp Asn Thr Asn Leu Pro Ser Glu Arg Ser Met Thr Gln Asn Gly 290 295 300			913
20	tat gcc gcc att gaa tag aactgttggt gcagcgttta cgccttttat Tyr Ala Ala Ile Glu 305			961
25	atgttattct ggtggtacct gtaaccatat aacgttgcat ttcotgtggt tgtaccattt ctctgataga ttttgaata atttgaagge aaagatagat tattgtgtag agagctacaa atttaattgat aaattgatca toagcactgg aaagctaaaa aaa			1021 1081 1124
30	<210> 6 <211> 309 <212> PRT <213> Arabidopsis thaliana			
35	Met Glu Met Ser Leu Arg Leu Ala Ser Ser Ser Thr Ser Asn Pro Ile 1 5 10 15			
40	Cys Leu Leu Asn Pro Gly Lys Asn Leu Asn Phe Pro Ile Arg Asn His 20 25 30			
45	Arg Ile Pro Lys Thr Ser Lys Pro Phe Cys Val Arg Ser Ser Met Ser 35 40 45			
50	Leu Ser Lys Pro Pro Arg Gln Thr Leu Ser Ser Asn Trp Asp Val Ser 50 55 60			
55	Ser Phe Ser Ile Asp Ser Val Ala Gln Ser Pro Ser Arg Leu Pro Ser 65 70 75 80			
60	Phe Glu Glu Leu Asp Thr Thr Asn Met Leu Leu Arg Gln Arg Ile Val 85 90 95			
	Phe Leu Gly Ser Gln Val Asp Asp Met Thr Ala Asp Leu Val Ile Ser 100 105 110			
	Gln Leu Leu Leu Leu Asp Ala Glu Asp Ser Glu Arg Asp Ile Thr Leu 115 120 125			

8

Phe Ile Asn Ser Pro Gly Gly Ser Ile Thr Ala Gly Met Gly Ile Tyr
 130 135 140

5 Asp Ala Met Lys Gln Cys Lys Ala Asp Val Ser Thr Val Cys Leu Gly
 145 150 155 160

10 Leu Ala Ala Ser Met Gly Ala Phe Leu Leu Ala Ser Gly Ser Lys Gly
 165 170 175

15 Lys Arg Tyr Cys Met Pro Asn Ser Lys Val Met Ile His Gln Pro Leu
 180 185 190

20 Gly Thr Ala Gly Gly Lys Ala Thr Glu Met Ser Ile Arg Ile Arg Glu
 195 200 205

Met Met Tyr His Lys Ile Lys Leu Asn Lys Ile Phe Ser Arg Ile Thr
 210 215 220

25 Gly Lys Pro Glu Ser Glu Ile Glu Ser Asp Thr Asp Arg Asp Asn Phe
 225 230 235 240

30 Leu Asn Pro Trp Glu Ala Lys Glu Tyr Gly Leu Ile Asp Ala Val Ile
 245 250 255

35 Asp Asp Gly Lys Pro Gly Leu Ile Ala Pro Ile Gly Asp Gly Thr Pro
 260 265 270

40 Pro Pro Lys Thr Lys Val Trp Asp Leu Trp Lys Val Glu Gly Thr Lys
 275 280 285

Lys Asp Asn Thr Asn Leu Pro Ser Glu Arg Ser Met Thr Gln Asn Gly
 290 295 300

45 Tyr Ala Ala Ile Glu
 305

50 <210> 7
 <211> 1183
 <212> DNA
 <213> Arabidopsis thaliana

55 <220>
 <221> CDS
 <222> (3) .. (902)
 <223>

60 <400> 7
 ct ttc ttc ttc ttc get toa gcc atg gga acc cta tct ctc tca tct
 Phe Phe Phe Phe Ala Ser Ala Met Gly Thr Leu Ser Leu Ser Ser
 1 5 10 15

47

9

	tct ctc aaa cct tca ctc gtt tca tca aga ctc aat tca tct tcc tcc	95
	Ser Leu Lys Pro Ser Leu Val Ser Ser Arg Leu Asn Ser Ser Ser Ser	
	20 25 30	
5	gca tct tct tct tct tct tct tct tct tct tct tct tct tct tct tct	143
	Ala Ser Ser Ser Ser Phe Pro Lys Pro Asn Asn Leu Tyr Leu Lys Pro	
	35 40 45	
10	acc aaa ctc att tca cca cct ctc aga aca act tca cca tct cca ttg	191
	Thr Lys Leu Ile Ser Pro Pro Leu Arg Thr Thr Ser Pro Ser Pro Leu	
	50 55 60	
15	aga ttc gcc aat gct tca atc gag atg tct cag aca cag gaa tca gct	239
	Arg Phe Ala Asn Ala Ser Ile Glu Met Ser Gln Thr Gln Glu Ser Ala	
	65 70 75	
20	att cgc gga gct gaa tct gac gtc atg ggt ctt ctc ctt agg gaa cga	287
	Ile Arg Gly Ala Glu Ser Asp Val Met Gly Leu Leu Leu Arg Glu Arg	
	80 85 90 95	
	atc gtc ttt ctc ggt agt agt atc gac gat ttc gtc gct gat gct att	335
	Ile Val Phe Leu Gly Ser Ser Ile Asp Asp Phe Val Ala Asp Ala Ile	
	100 105 110	
25	atg agt cag ttg ctt ctc tta gat gct aaa gat cca aag aaa gat atc	383
	Met Ser Gln Leu Leu Leu Leu Asp Ala Lys Asp Pro Lys Lys Asp Ile	
	115 120 125	
30	aaa ctc ttt atc aat cct cct ggt ggt tct ctc agt gca acc atg gct	431
	Lys Leu Phe Ile Asn Ser Pro Gly Gly Ser Leu Ser Ala Thr Met Ala	
	130 135 140	
35	ata tac gat gtc gtt cag ctt gtg aga gct gat gtt tct acg att gct	479
	Ile Tyr Asp Val Val Gln Leu Val Arg Ala Asp Val Ser Thr Ile Ala	
	145 150 155	
40	ctt ggc att got gca tca aca gct tct att att ctt ggt gtc gga act	527
	Leu Gly Ile Ala Ala Ser Thr Ala Ser Ile Ile Leu Gly Ala Gly Thr	
	160 165 170 175	
	aaa ggc aag cgc ttt gct atg ccc aac acg agg ata atg att cat caa	575
	Lys Gly Lys Arg Phe Ala Met Pro Asn Thr Arg Ile Met Ile His Gln	
	180 185 190	
45	cct ctt gga ggt gca agc ggt caa got ata gat gtt gag att caa gct	623
	Pro Leu Gly Gly Ala Ser Gly Gln Ala Ile Asp Val Glu Ile Gln Ala	
	195 200 205	
50	aag gaa gtt atg cat aac aag aac aat gtc acc agc att atc gcg gga	671
	Lys Glu Val Met His Asn Lys Asn Asn Val Thr Ser Ile Ile Ala Gly	
	210 215 220	
55	tgt act agt cga tct ttt gag cag gtt ctg aaa gat att gat agg gac	719
	Cys Thr Ser Arg Ser Phe Glu Gln Val Leu Lys Asp Ile Asp Arg Asp	
	225 230 235	
60	cgg tac atg tct cca att gaa gca gtt gag tat ggt tta att gat gga	767
	Arg Tyr Met Ser Pro Ile Glu Ala Val Glu Tyr Gly Leu Ile Asp Gly	
	240 245 250 255	
	gtt att gat gga gac agc att att cct ctt gaa cot gtt cct gat aga	815
	Val Ile Asp Gly Asp Ser Ile Ile Pro Leu Glu Pro Val Pro Asp Arg	
	260 265 270	

10

	gtg aaa cgg aga gta aac tac gag gag att agc aag gat cgg atg aaa	863
	Val Lys Pro Arg Val Asn Tyr Glu Glu Ile Ser Lys Asp Pro Met Lys	
	275 280 285	
5	ttc ttg act ccc gag ata cct gat gat gag atc tac taa agccaagctc	912
	Phe Leu Thr Pro Glu Ile Pro Asp Asp Glu Ile Tyr	
	290 295	
10	gtctagaagc agggatcttc aaatgtgact aagactagca gtttcgagga aaagctcaat	972
	ttctttctgcg gttactggta ttggctttgc gaaaccgaag ctggtagtac ttggctttcg	1032
	tatctcatat ttcagttgtt cagaaaataa ttgttcttta aatcactctg ttttgaggaa	1092
15	aatgacttaa agaagctgta gttatctcgt ttatgacaat cccttcaagt gtttaaatgga	1152
	ttcaagaagt atcagtcagt atttttctgtg t	1163
20	<210> 8	
	<211> 299	
	<212> PRT	
	<213> Arabidopsis thaliana	
25	<400> 8	
	Phe Phe Phe Phe Ala Ser Ala Met Gly Thr Leu Ser Leu Ser Ser Ser	
	1 5 10 15	
30	Leu Lys Pro Ser Leu Val Ser Ser Arg Leu Asn Ser Ser Ser Ser Ala	
	20 25 30	
35	Ser Ser Ser Ser Phe Pro Lys Pro Asn Asn Leu Tyr Leu Lys Pro Thr	
	35 40 45	
40	Lys Leu Ile Ser Pro Pro Leu Arg Thr Thr Ser Pro Ser Pro Leu Arg	
	50 55 60	
45	Phe Ala Asn Ala Ser Ile Glu Met Ser Gln Thr Gln Glu Ser Ala Ile	
	65 70 75 80	
	Arg Gly Ala Glu Ser Asp Val Met Gly Leu Leu Leu Arg Glu Arg Ile	
	85 90 95	
50	Val Phe Leu Gly Ser Ser Ile Asp Asp Phe Val Ala Asp Ala Ile Met	
	100 105 110	
55	Ser Gln Leu Leu Leu Leu Asp Ala Lys Asp Pro Lys Lys Asp Ile Lys	
	115 120 125	
60	Leu Phe Ile Asn Ser Pro Gly Gly Ser Leu Ser Ala Thr Met Ala Ile	
	130 135 140	
	Tyr Asp Val Val Gln Leu Val Arg Ala Asp Val Ser Thr Ile Ala Leu	
	145 150 155 160	

5	Gly Ile Ala Ala Ser Thr Ala Ser Ile Ile Leu Gly Ala Gly Thr Lys 165 170 175	
10	Gly Lys Arg Phe Ala Met Pro Asn Thr Arg Ile Met Ile His Gln Pro 180 185 190	
15	Leu Gly Gly Ala Ser Gly Gln Ala Ile Asp Val Glu Ile Gln Ala Lys 195 200 205	
20	Glu Val Met His Asn Lys Asn Asn Val Thr Ser Ile Ile Ala Gly Cys 210 215 220	
25	Thr Ser Arg Ser Phe Glu Gln Val Leu Lys Asp Ile Asp Arg Asp Arg 225 230 235 240	
30	Tyr Met Ser Pro Ile Glu Ala Val Glu Tyr Gly Leu Ile Asp Gly Val 245 250 255	
35	Ile Asp Gly Asp Ser Ile Ile Pro Leu Glu Pro Val Pro Asp Arg Val 260 265 270	
40	Lys Pro Arg Val Asn Tyr Glu Glu Ile Ser Lys Asp Pro Met Lys Phe 275 280 285	
45	Leu Thr Pro Glu Ile Pro Asp Asp Glu Ile Tyr 290 295	
50	<210> 9 <211> 1056 <212> DNA <213> Arabidopsis thaliana	
55	<220> <221> CDS <222> (61)..(876) <223>	
60	<400> 9 gagtaattta gcatctatcc acgcttgaac ccgaaaaact ctgaaagctg agctctggta atg gcg ggt tta gca att tca cct cct ctc ggt ctt tcc ttc tct tct Met Ala Gly Leu Ala Ile Ser Pro Pro Leu Gly Leu Ser Phe Ser Ser 1 5 10 15 cga act cga aac cct aaa ccc act tcc ttt cta tct cac aat caa agg Arg Thr Arg Asn Pro Lys Pro Thr Ser Phe Leu Ser His Asn Gln Arg 20 25 30 aat cct ata aga cgt ata gtt tct gct cta cag agt cca tat gga gat Asn Pro Ile Arg Arg Ile Val Ser Ala Leu Gln Ser Pro Tyr Gly Asp 35 40 45	60 108 156 204

12

	tct ctg aaa gct gga ctt tct agt aat gtt tct gga tcc cca ata aag Ser Leu Lys Ala Gly Leu Ser Ser Asn Val Ser Gly Ser Pro Ile Lys 50 55 60	252
5	att gac aac aag gct cca aga ttt gga gtg ata gag gcg aaa aag gga Ile Asp Asn Lys Ala Pro Arg Phe Gly Val Ile Glu Ala Lys Lys Gly 65 70 75 80	300
10	aac ccc cca gta atg cct tca gtg atg acc cct gga gga cct tta gac Asn Pro Pro Val Met Pro Ser Val Met Thr Pro Gly Gly Pro Leu Asp 85 90 95	348
15	ctc tct tct gtg tta ttc cgt aac cgc ata atc ttc atc ggg caa cca Leu Ser Ser Val Leu Phe Arg Asn Arg Ile Ile Phe Ile Gly Gln Pro 100 105 110	396
20	att aac gca cag gtt gct cag cga gtc ata tct cag ctt gta acc ctt Ile Asn Ala Gln Val Ala Gln Arg Val Ile Ser Gln Leu Val Thr Leu 115 120 125	444
	gca tct att gat gat aaa tcc gac atc ctg atg tac ttg aat tgt ccc Ala Ser Ile Asp Asp Lys Ser Asp Ile Leu Met Tyr Leu Asn Cys Pro 130 135 140	492
25	ggg ggc agt act tac tcc gtc cta aca att tat gac tgt atg tct tgg Gly Gly Ser Thr Tyr Ser Val Leu Thr Ile Tyr Asp Cys Met Ser Trp 145 150 155 160	540
30	ata aag cct aaa gtt gga aca gtg gcg ttt gga gta gct gca agc caa Ile Lys Pro Lys Val Gly Thr Val Ala Phe Gly Val Ala Ala Ser Gln 165 170 175	588
35	gga gca ttt ttt ctt gct gga ggt gaa aaa gga atg cgt tat gca atg Gly Ala Phe Phe Leu Ala Gly Gly Glu Lys Gly Met Arg Tyr Ala Met 180 185 190	636
40	cca aat act cgt gtc atg ata cat caa cca caa act gga tgc gga gga Pro Asn Thr Arg Val Met Ile His Gln Pro Gln Thr Gly Cys Gly Gly 195 200 205	684
	cat gta gag gac gtg agg aga cag gtc aat gaa gcc atc gaa gcc cga His Val Glu Asp Val Arg Arg Gln Val Asn Glu Ala Ile Glu Ala Arg 210 215 220	732
45	caa aaa att gac agg atg tat gca gct ttc act gga caa cct ctg gag Gln Lys Ile Asp Arg Met Tyr Ala Ala Phe Thr Gly Gln Pro Leu Glu 225 230 235 240	780
50	aaa gtg cag caa tac act gaa aga gat cgt ttc tta tca gca tct gag Lys Val Gln Gln Tyr Thr Glu Arg Asp Arg Phe Leu Ser Ala Ser Glu 245 250 255	828
55	gcg ttt gag ttc ggg ctc att gat ggt cta ttg gaa aca gaa tac tga Ala Phe Glu Phe Gly Leu Ile Asp Gly Leu Leu Glu Thr Glu Tyr 260 265 270	876
	agcagcatac aggacaatgc acaacaacag ctcatctgcaa tggttcaaagc ttccattttc	936
60	atattgaatat gaacggttgt aactgatatt tgtgcataaa tcagtttgggt tttcttgggt	996
	ttattgtota ctaaaacagaa tgagaaaact aaactgttta ttttttttact gaaaaatctg	1056

<210> 10

13

<211> 271
<212> PRT
<213> Arabidopsis thaliana

5 <400> 10

Met Ala Gly Leu Ala Ile Ser Pro Pro Leu Gly Leu Ser Phe Ser Ser
1 5 10 15

10

Arg Thr Arg Asn Pro Lys Pro Thr Ser Phe Leu Ser His Asn Gln Arg
20 25 30

15

Asn Pro Ile Arg Arg Ile Val Ser Ala Leu Gln Ser Pro Tyr Gly Asp
35 40 45

20

Ser Leu Lys Ala Gly Leu Ser Ser Asn Val Ser Gly Ser Pro Ile Lys
50 55 60

25

Ile Asp Asn Lys Ala Pro Arg Phe Gly Val Ile Glu Ala Lys Lys Gly
65 70 75 80

30

Asn Pro Pro Val Met Pro Ser Val Met Thr Pro Gly Gly Pro Leu Asp
85 90 95

35

Leu Ser Ser Val Leu Phe Arg Asn Arg Ile Ile Phe Ile Gly Gln Pro
100 105 110

40

Ile Asn Ala Gln Val Ala Gln Arg Val Ile Ser Gln Leu Val Thr Leu
115 120 125

45

Ala Ser Ile Asp Asp Lys Ser Asp Ile Leu Met Tyr Leu Asn Cys Pro
130 135 140

Gly Gly Ser Thr Tyr Ser Val Leu Thr Ile Tyr Asp Cys Met Ser Trp
145 150 155 160

50

Ile Lys Pro Lys Val Gly Thr Val Ala Phe Gly Val Ala Ala Ser Gln
165 170 175

55

Gly Ala Phe Phe Leu Ala Gly Gly Glu Lys Gly Met Arg Tyr Ala Met
180 185 190

60

Pro Asn Thr Arg Val Met Ile His Gln Pro Gln Thr Gly Cys Gly Gly
195 200 205

His Val Glu Asp Val Arg Arg Gln Val Asn Glu Ala Ile Glu Ala Arg
210 215 220

Gln Lys Ile Asp Arg Met Tyr Ala Ala Phe Thr Gly Gln Pro Leu Glu
225 230 235 240

14

5	Lys Val Gln Gln Tyr Thr Glu Arg Asp Arg Phe Leu Ser Ala Ser Glu	245	250	255
10	Ala Phe Glu Phe Gly Leu Ile Asp Gly Leu Leu Glu Thr Glu Tyr	260	265	270
15	<210> 11 <211> 1448 <212> DNA <213> Nicotiana tabacum			
20	<220> <221> CDS <222> (2)..(1162) <223>			
25	<400> 11 g cgg ccg ctg gct tct tct ttg ctt ctc tct ccg ctt tct agc tcg acg Arg Pro Leu Ala Ser Ser Leu Leu Leu Ser Pro Leu Ser Ser Ser Thr	1	5	10 15
30	gtt act gaa aat cgc gag ctg ggt tct ggt aaa tca act ttc ata tcc Val Thr Glu Asn Arg Glu Leu Gly Ser Gly Lys Ser Thr Phe Ile Ser	20	25	30
35	agt ccc aat ttc tcc ttt gca act tct gtt cac agt tgc agg cca aac Ser Pro Asn Phe Ser Phe Ala Thr Ser Val His Ser Cys Arg Pro Asn	35	40	45
40	ggc gtt cga ggt tat tgt tac agg tct cgg gta gct aag tct ttg gac Gly Val Arg Gly Tyr Cys Tyr Arg Ser Pro Val Ala Lys Ser Leu Asp	50	55	60
45	cat ata ccc caa aaa ttc aga ctg gaa aat ctc aaa gat gga cta ctg His Ile Pro Gln Lys Phe Arg Leu Glu Asn Leu Lys Asp Gly Leu Leu	65	70	75 80
50	gac aac tat aaa agt gcc cct cag tat ctt tac ggc ctt agt cct tca Asp Asn Tyr Lys Ser Ala Pro Gln Tyr Leu Tyr Gly Leu Ser Pro Ser	85	90	95
55	cag atg gat atg ttc atg aca gaa gat aac cca gta cgg cga cag tca Gln Met Asp Met Phe Met Thr Glu Asp Asn Pro Val Arg Arg Gln Ser	100	105	110
60	gaa agt gcc act gag gat agt ata tct tca gcc aat aac tat ctg agc Glu Ser Ala Thr Glu Asp Ser Ile Ser Ser Ala Asn Asn Tyr Leu Ser	115	120	125
65	aat ggt gga atg tgg agt atg tcc ggc atg aac gat cgg ggc ccc tcg Asn Gly Gly Met Trp Ser Met Ser Gly Met Asn Asp Arg Gly Pro Ser	130	135	140
70	aaa tac agt atg agt gtc agc atg tac cgt gga gga aca aga gga tct Lys Tyr Ser Met Ser Val Ser Met Tyr Arg Gly Gly Thr Arg Gly Ser	145	150	155 160
75	gga aga cct cga act gcg cct cct gat ttg cca tct ttg ctt ttg gat Gly Arg Pro Arg Thr Ala Pro Pro Asp Leu Pro Ser Leu Leu Leu Asp			

15

	165	170	175	
5	get cga att gtc tat ctg ggc atg cct att gta cca gct gtt aca gag Ala Arg Ile Val Tyr Leu Gly Met Pro Ile Val Pro Ala Val Thr Glu 180 185 190	577		
10	ctt ctt gtt get cag ttt atg tgg ttg gat tat gac aat cca tca aag Leu Leu Val Ala Gln Phe Met Trp Leu Asp Tyr Asp Asn Pro Ser Lys 195 200 205	625		
15	cct ata tac cta tat ata aac tca tca ggc aca cag aat gag aag atg Pro Ile Tyr Leu Tyr Ile Asn Ser Ser Gly Thr Gln Asn Glu Lys Met 210 215 220	673		
20	gag act gtt ggg tct gaa aca gag gca tat gcc atc gct gac aca atg Glu Thr Val Gly Ser Glu Thr Glu Ala Tyr Ala Ile Ala Asp Thr Met 225 230 235 240	721		
25	gca tac tgc aaa tca gat atc tat aca gtg aac tgt ggc atg goa tat Ala Tyr Cys Lys Ser Asp Ile Tyr Thr Val Asn Cys Gly Met Ala Tyr 245 250 255	769		
30	ggc caa gca gca atg ctt ctg tca ctg gga aag aag ggg ttc cgt gct Gly Gln Ala Ala Met Leu Leu Ser Leu Gly Lys Lys Gly Phe Arg Ala 260 265 270	817		
35	atg cag cca aat tca tct aca aaa ttg tat cta cct aaa gtc agc aaa Met Gln Pro Asn Ser Ser Thr Lys Leu Tyr Leu Pro Lys Val Ser Lys 275 280 285	865		
40	tcc agt gga gca gtg ata gat atg tgg atc agg gcc aaa gaa cta gat Ser Ser Gly Ala Val Ile Asp Met Trp Ile Arg Ala Lys Glu Leu Asp 290 295 300	913		
45	gca aac act gag tat tac ctt gaa cta tta gcg aaa gga gtt gga aaa Ala Asn Thr Glu Tyr Tyr Leu Glu Leu Leu Ala Lys Gly Val Gly Lys 305 310 315 320	961		
50	cca aag gaa gaa atc gag aaa gat att caa cgc cct aaa tat ctg cgg Pro Lys Glu Glu Ile Glu Lys Asp Ile Gln Arg Pro Lys Tyr Leu Arg 325 330 335	1009		
55	gca caa gaa gcc att gac tat ggc att gcg gac aag ata atc gat tca Ala Gln Glu Ala Ile Asp Tyr Gly Ile Ala Asp Lys Ile Ile Asp Ser 340 345 350	1057		
60	aga gac aat gca ttt gag aaa agg aac tat ggt gag ata ctc gcc caa Arg Asp Asn Ala Phe Glu Lys Arg Asn Tyr Gly Glu Ile Leu Ala Gln 355 360 365	1105		
65	tct aga gct atg agg aaa gcc gga cca ggt gct cag gct gct cca tct Ser Arg Ala Met Arg Lys Ala Gly Pro Gly Ala Gln Ala Ala Pro Ser 370 375 380	1153		
70	ggc tcc agg tgactggaag agcggtaatg gtcccaagct ttcaggaaca Gly Ser Arg 385	1202		
75	actgtctgttc ccttatagtt tcgaggaaca aagttgctgg ttacttggtc tgtgccggtc taagtgaact gggacaaaga acatattgta gaaacottgt ttgagctgtg aagtataggg gtttttacaac tattatgcac aggtctgcaa agagtaccca taatgtcaat tggttgtacc	1262 1322 1382		

16

agtatcaaac aatcagatag tgccagtgtg ttggtataaat gaatatagat ctctctgagc 1442

ggccgc 1448

5

<210> 12

<211> 387

<212> PRT

<213> Nicotiana tabacum

10

<400> 12

15

Arg Pro Leu Ala Ser Ser Leu Leu Leu Ser Pro Leu Ser Ser Ser Thr
1 5 10 15Val Thr Glu Asn Arg Glu Leu Gly Ser Gly Lys Ser Thr Phe Ile Ser
20 25 30

20

Ser Pro Asn Phe Ser Phe Ala Thr Ser Val His Ser Cys Arg Pro Asn
35 40 45

25

Gly Val Arg Gly Tyr Cys Tyr Arg Ser Pro Val Ala Lys Ser Leu Asp
50 55 60

30

His Ile Pro Gln Lys Phe Arg Leu Glu Asn Leu Lys Asp Gly Leu Leu
65 70 75 80

35

Asp Asn Tyr Lys Ser Ala Pro Gln Tyr Leu Tyr Gly Leu Ser Pro Ser
85 90 95

40

Gln Met Asp Met Phe Met Thr Glu Asp Asn Pro Val Arg Arg Gln Ser
100 105 110Glu Ser Ala Thr Glu Asp Ser Ile Ser Ser Ala Asn Asn Tyr Leu Ser
115 120 125

45

Asn Gly Gly Met Trp Ser Met Ser Gly Met Asn Asp Arg Gly Pro Ser
130 135 140

50

Lys Tyr Ser Met Ser Val Ser Met Tyr Arg Gly Gly Thr Arg Gly Ser
145 150 155 160

55

Gly Arg Pro Arg Thr Ala Pro Pro Asp Leu Pro Ser Leu Leu Leu Asp
165 170 175Ala Arg Ile Val Tyr Leu Gly Met Pro Ile Val Pro Ala Val Thr Glu
180 185 190

60

Leu Leu Val Ala Gln Phe Met Trp Leu Asp Tyr Asp Asn Pro Ser Lys
195 200 205

17

Pro Ile Tyr Leu Tyr Ile Asn Ser Ser Gly Thr Gln Asn Glu Lys Met
 210 215 220

5 Glu Thr Val Gly Ser Glu Thr Glu Ala Tyr Ala Ile Ala Asp Thr Met
 225 230 235 240

10 Ala Tyr Cys Lys Ser Asp Ile Tyr Thr Val Asn Cys Gly Met Ala Tyr
 245 250 255

15 Gly Gln Ala Ala Met Leu Leu Ser Leu Gly Lys Lys Gly Phe Arg Ala
 260 265 270

20 Met Gln Pro Asn Ser Ser Thr Lys Leu Tyr Leu Pro Lys Val Ser Lys
 275 280 285

Ser Ser Gly Ala Val Ile Asp Met Trp Ile Arg Ala Lys Glu Leu Asp
 290 295 300

25 Ala Asn Thr Glu Tyr Tyr Leu Glu Leu Leu Ala Lys Gly Val Gly Lys
 305 310 315 320

30 Pro Lys Glu Glu Ile Glu Lys Asp Ile Gln Arg Pro Lys Tyr Leu Arg
 325 330 335

35 Ala Gln Glu Ala Ile Asp Tyr Gly Ile Ala Asp Lys Ile Ile Asp Ser
 340 345 350

Arg Asp Asn Ala Phe Glu Lys Arg Asn Tyr Gly Glu Ile Leu Ala Gln
 355 360 365

40 Ser Arg Ala Met Arg Lys Ala Gly Pro Gly Ala Gln Ala Ala Pro Ser
 370 375 380

45 Gly Ser Arg
 385

50 <210> 13
 <211> 1246
 <212> DNA
 <213> Arabidopsis thaliana

55 <220>
 <221> CDS
 <222> (38)..(1030)
 <223>

60 <400> 13
 atttttcgcga gcttccgtgt ccaagagctc ctogacc atg gcg tct tgt tta caa 55
 Met Ala Ser Cys Leu Gln
 1 5

18

	gca tcc atg aat tct ctg ctt cca cgc tct tct tct ttt tct cct cat	103
	Ala Ser Met Asn Ser Leu Leu Pro Arg Ser Ser Ser Phe Ser Pro His	
	10 15 20	
5	cct cct cta tct tct aat tca tcc ggg aga aga aac ttg aag act ttt	151
	Pro Pro Leu Ser Ser Asn Ser Ser Gly Arg Arg Asn Leu Lys Thr Phe	
	25 30 35	
10	cgt tac gcc ttt cgc gcc aaa gcc tct gcc aaa atc cct atg cct cgg	199
	Arg Tyr Ala Phe Arg Ala Lys Ala Ser Ala Lys Ile Pro Met Pro Pro	
	40 45 50	
15	ata aat cca aag gat cct ttc ctc tcc acg ctc gct tct att gcc gcg	247
	Ile Asn Pro Lys Asp Pro Phe Leu Ser Thr Leu Ala Ser Ile Ala Ala	
	55 60 65 70	
20	aat tct ccg gaa aag ctt ctc aat cgg ccg gtt aac gct gat gtg ccg	295
	Asn Ser Pro Glu Lys Leu Leu Asn Arg Pro Val Asn Ala Asp Val Pro	
	75 80 85	
25	cca tat ctt gac atc ttt gac tcc cct cag ctc atg tct tct cct gca	343
	Pro Tyr Leu Asp Ile Phe Asp Ser Pro Gln Leu Met Ser Ser Pro Ala	
	90 95 100	
30	cag gtt gaa aga tca gtg gct tat aac gag cac cga ccg aga act cct	391
	Gln Val Glu Arg Ser Val Ala Tyr Asn Glu His Arg Pro Arg Thr Pro	
	105 110 115	
35	cca cca gac ttg cca tct atg ctt ctt gac ggg aga att gtt tac att	439
	Pro Pro Asp Leu Pro Ser Met Leu Leu Asp Gly Arg Ile Val Tyr Ile	
	120 125 130	
40	gga atg cct ctt gtg ccg gca gtg act gag cta gtt gtc gct gag cta	487
	Gly Met Pro Leu Val Pro Ala Val Thr Glu Leu Val Val Ala Glu Leu	
	135 140 145 150	
45	atg tat ctt cag tgg ctg gat ccc aag gaa ccc att tac att tac atc	535
	Met Tyr Leu Gln Trp Leu Asp Pro Lys Glu Pro Ile Tyr Ile Tyr Ile	
	155 160 165	
50	aac tcc aca ggg acc act cgt gat gat gga gag acg gtt gga atg gaa	583
	Asn Ser Thr Gly Thr Thr Arg Asp Asp Gly Glu Thr Val Gly Met Glu	
	170 175 180	
55	tca gaa ggg ttt gcg atc tat gac tct ttg atg caa ctt aaa aac gag	631
	Ser Glu Gly Phe Ala Ile Tyr Asp Ser Leu Met Gln Lys Asn Glu	
	185 190 195	
60	gta cat aca gta tgt gtg gga gca gcc ata ggt cag gcc tgt cta tta	679
	Val His Thr Val Cys Val Gly Ala Ala Ile Gly Gln Ala Cys Leu Leu	
	200 205 210	
65	ctt tct gcg gga acc aag ggt aaa cgg ttt atg atg cca cat gcc aaa	727
	Leu Ser Ala Gly Thr Lys Gly Lys Arg Phe Met Met Pro His Ala Lys	
	215 220 225 230	
70	gcg atg att cag caa coa cgt gta cct tct tct ggg ttg atg cct gcc	775
	Ala Met Ile Gln Gln Pro Arg Val Pro Ser Ser Gly Leu Met Pro Ala	
	235 240 245	
75	agt gat gtc ctg att cgg gcc aaa gag gtc att aca aat agg gat ata	823
	Ser Asp Val Leu Ile Arg Ala Lys Glu Val Ile Thr Asn Arg Asp Ile	
	250 255 260	

19

	ctt	gtg	gaa	cta	cta	tca	aag	cat	act	ggg	aat	tcc	gtg	gag	act	gta	871
	Leu	Val	Glu	Leu	Leu	Ser	Lys	His	Thr	Gly	Asn	Ser	Val	Glu	Thr	Val	
			265					270					275				
5	gct	aac	gta	atg	aga	agg	cca	tat	tac	atg	gat	gca	cca	aaa	gct	aaa	919
	Ala	Asn	Val	Met	Arg	Arg	Pro	Tyr	Tyr	Met	Asp	Ala	Pro	Lys	Ala	Lys	
			280				285					290					
10	gaa	ttt	gga	gtc	att	gac	agg	att	ctt	tgg	cgc	ggg	caa	gaa	aag	att	967
	Glu	Phe	Gly	Val	Ile	Asp	Arg	Ile	Leu	Trp	Arg	Gly	Gln	Glu	Lys	Ile	
						300					305					310	
15	att	gcg	gac	gtg	gtt	oct	tca	gag	gaa	ttc	gac	aag	aat	gca	ggg	att	1015
	Ile	Ala	Asp	Val	Val	Pro	Ser	Glu	Glu	Phe	Asp	Lys	Asn	Ala	Gly	Ile	
					315					320					325		
20	aaa	agc	gta	gta	tga	gtctagtcctt	aagttttctt	ggcctaaatc	atactgcgtc								1070
	Lys	Ser	Val	Val													
				330													
	atggagaaga	acaaatagac	tgacccaaat	cacattggcc	gcagactgcc	ttgtttcaaa											1130
	tcacttggtta	aatgtgaaca	tgcgattagg	agaatcatac	ttaaaggatc	ttgaaatatt											1190
25	atgataaaat	tgtaatgtgt	ttgttcgtta	gcaatagtaa	atacaatctt	caactc											1246
	<210>	14															
	<211>	330															
30	<212>	PRT															
	<213>	Arabidopsis thaliana															
	<400>	14															
35	Met	Ala	Ser	Cys	Leu	Gln	Ala	Ser	Met	Asn	Ser	Leu	Leu	Pro	Arg	Ser	
	1				5					10					15		
40	Ser	Ser	Phe	Ser	Pro	His	Pro	Pro	Leu	Ser	Ser	Asn	Ser	Ser	Gly	Arg	
				20					25						30		
45	Arg	Asn	Leu	Lys	Thr	Phe	Arg	Tyr	Ala	Phe	Arg	Ala	Lys	Ala	Ser	Ala	
			35					40					45				
50	Lys	Ile	Pro	Met	Pro	Pro	Ile	Asn	Pro	Lys	Asp	Pro	Phe	Leu	Ser	Thr	
		50					55					60					
55	Leu	Ala	Ser	Ile	Ala	Ala	Asn	Ser	Pro	Glu	Lys	Leu	Leu	Asn	Arg	Pro	
	65					70					75				80		
60	Val	Asn	Ala	Asp	Val	Pro	Pro	Tyr	Leu	Asp	Ile	Phe	Asp	Ser	Pro	Gln	
				85					90						95		
65	Leu	Met	Ser	Ser	Pro	Ala	Gln	Val	Glu	Arg	Ser	Val	Ala	Tyr	Asn	Glu	
				100					105					110			
70	His	Arg	Pro	Arg	Thr	Pro	Pro	Pro	Asp	Leu	Pro	Ser	Met	Leu	Leu	Asp	
				115				120					125				

20

- 5 Gly Arg Ile Val Tyr Ile Gly Met Pro Leu Val Pro Ala Val Thr Glu
130 135 140
- 10 Leu Val Val Ala Glu Leu Met Tyr Leu Gln Trp Leu Asp Pro Lys Glu
145 150 155 160
- 15 Pro Ile Tyr Ile Tyr Ile Asn Ser Thr Gly Thr Thr Arg Asp Asp Gly
165 170 175
- 20 Glu Thr Val Gly Met Glu Ser Glu Gly Phe Ala Ile Tyr Asp Ser Leu
180 185 190
- 25 Met Gln Leu Lys Asn Glu Val His Thr Val Cys Val Gly Ala Ala Ile
195 200 205
- 30 Gly Gln Ala Cys Leu Leu Leu Ser Ala Gly Thr Lys Gly Lys Arg Phe
210 215 220
- 35 Met Met Pro His Ala Lys Ala Met Ile Gln Gln Pro Arg Val Pro Ser
225 230 235 240
- 40 Ser Gly Leu Met Pro Ala Ser Asp Val Leu Ile Arg Ala Lys Glu Val
245 250 255
- 45 Ile Thr Asn Arg Asp Ile Leu Val Glu Leu Leu Ser Lys His Thr Gly
260 265 270
- 50 Asn Ser Val Glu Thr Val Ala Asn Val Met Arg Arg Pro Tyr Tyr Met
275 280 285
- 55 Asp Ala Pro Lys Ala Lys Glu Phe Gly Val Ile Asp Arg Ile Leu Trp
290 295 300
- 60 Arg Gly Gln Glu Lys Ile Ile Ala Asp Val Val Pro Ser Glu Glu Phe
305 310 315 320
- 65 Asp Lys Asn Ala Gly Ile Lys Ser Val Val
325 330
- 70 <210> 15
<211> 1236
<212> DNA
<213> Arabidopsis thaliana
- 75 <220>
<221> CDS
<222> (66)..(983)
<223>

	<400> 15	
	agatcggttat cgttttcgggg tcacagggac tctcactctt tctctctctc tgcacacaaag	60
5	aagaa atg gag gta gca gca gcg act gcg acg agc ttc aca acg ott cga Met Glu Val Ala Ala Ala Thr Ala Thr Ser Phe Thr Thr Leu Arg 1 5 10 15	110
10	gct cgt acg tca gcg att atc ccg tct tct aca cgt aat ctg aga tct Ala Arg Thr Ser Ala Ile Ile Pro Ser Ser Thr Arg Asn Leu Arg Ser 20 25 30	158
15	aaa ccg aga ttt tct tca tot tca tct ctg aga gct tct ctt tcg aat Lys Pro Arg Phe Ser Ser Ser Ser Ser Leu Arg Ala Ser Leu Ser Asn 35 40 45	206
20	ggc ttt ctt tcg ccg tat acc gga gga agc atc tct agt gac tta tgc Gly Phe Leu Ser Pro Tyr Thr Gly Gly Ser Ile Ser Ser Asp Leu Cys 50 55 60	254
	ggc gct aag ctt cgt gcg gaa tcg ott aat ccg tta aat ttt tcc agt Gly Ala Lys Leu Arg Ala Glu Ser Leu Asn Pro Leu Asn Phe Ser Ser 65 70 75	302
25	tcc aag cct aaa cgc gga gtt gtc act atg gtt ata cct ttc tca aag Ser Lys Pro Lys Arg Gly Val Val Thr Met Val Ile Pro Phe Ser Lys 80 85 90 95	350
30	gga agt gca cac gaa caa cct cct cct gat ttg gca tca tat ttg ttc Gly Ser Ala His Glu Gln Pro Pro Pro Asp Leu Ala Ser Tyr Leu Phe 100 105 110	398
35	aag aac cga att gta tat ttg gga atg tct ctg gta cct tca gtt act Lys Asn Arg Ile Val Tyr Leu Gly Met Ser Leu Val Pro Ser Val Thr 115 120 125	446
40	gag ttg ata ctt gcg gag ttt ctt tac ctt cag tat gaa gac gag gaa Glu Leu Ile Leu Ala Glu Phe Leu Tyr Leu Gln Tyr Glu Asp Glu Glu 130 135 140	494
	aag cct att tac ctt tac ata aac tcg act ggg aca acc aag aat ggt Lys Pro Ile Tyr Leu Tyr Ile Asn Ser Thr Gly Thr Thr Lys Asn Gly 145 150 155	542
45	gaa aag ttg ggc tat gat act gag gct ttt gca atc tat gat gtc atg Glu Lys Leu Gly Tyr Asp Thr Glu Ala Phe Ala Ile Tyr Asp Val Met 160 165 170 175	590
50	ggg tat gtc aaa cca cca atc ttt act ctt tgc gtc ggg aat gcg tgg Gly Tyr Val Lys Pro Pro Ile Phe Thr Leu Cys Val Gly Asn Ala Trp 180 185 190	638
55	ggt gaa gct gct ttg ctt ctg act gct ggt gca aaa gga aat cga tct Gly Glu Ala Ala Leu Leu Leu Thr Ala Gly Ala Lys Gly Asn Arg Ser 195 200 205	686
	gcg ttg ccc tca tca act att atg ata aag cag ccc att gct cga ttt Ala Leu Pro Ser Ser Thr Ile Met Ile Lys Gln Pro Ile Ala Arg Phe 210 215 220	734
60	caa ggc caa gca aot gat gtt gaa att gca agg aaa gaa atc aag cac Gln Gly Gln Ala Thr Asp Val Glu Ile Ala Arg Lys Glu Ile Lys His 225 230 235	782

22

	ata aag aca gaa atg gtc aag ctg tat tca aag cat att ggt aaa tcc	830
	Ile Lys Thr Glu Met Val Lys Leu Tyr Ser Lys His Ile Gly Lys Ser	
	240 245 250 255	
5	ccg gag cag att gaa gct gac arg aaa cgc ccg aaa tat ttt agt ccc	878
	Pro Glu Gln Ile Glu Ala Asp Met Lys Arg Pro Lys Tyr Phe Ser Pro	
	260 265 270	
10	act gag gct gtt gaa tat ggg atc att gat aag gtg gtt tac aat gaa	926
	Thr Glu Ala Val Glu Tyr Gly Ile Ile Asp Lys Val Val Tyr Asn Glu	
	275 280 285	
15	agg ggc agc caa gac aga gga gtt gtg tct gac ctt aaa aag gca caa	974
	Arg Gly Ser Gln Asp Arg Gly Val Val Ser Asp Leu Lys Lys Ala Gln	
	290 295 300	
	ctc att tga atgtcagaac tgtcttccga aatcccacga ttaacaggct	1023
	Leu Ile	
	305	
20	ggagatctta ccgctgacca aatgggggaat cagtgaacca ttcaccggca cagaactgag	1083
	gtaaagtctg gaaaacatgt taaaaaagggt tactagtaat gctgcaattg taggggtatt	1143
25	tgaacagaaa caaacccata tgtgtaggct tgtgaatgoc tagaaacagg attgggtgtat	1203
	cttcaatata tgtttctaag atgaatcaat ttc	1236
30	<210> 16	
	<211> 305	
	<212> PRT	
	<213> Arabidopsis thaliana	
35	<400> 16	
	Met Glu Val Ala Ala Ala Thr Ala Thr Ser Phe Thr Thr Leu Arg Ala	
	1 5 10 15	
40	Arg Thr Ser Ala Ile Ile Pro Ser Ser Thr Arg Asn Leu Arg Ser Lys	
	20 25 30	
45	Pro Arg Phe Ser Ser Ser Ser Ser Leu Arg Ala Ser Leu Ser Asn Gly	
	35 40 45	
50	Phe Leu Ser Pro Tyr Thr Gly Gly Ser Ile Ser Ser Asp Leu Cys Gly	
	50 55 60	
55	Ala Lys Leu Arg Ala Glu Ser Leu Asn Pro Leu Asn Phe Ser Ser Ser	
	65 70 75 80	
	Lys Pro Lys Arg Gly Val Val Thr Met Val Ile Pro Phe Ser Lys Gly	
	85 90 95	
60	Ser Ala His Glu Gln Pro Pro Pro Asp Leu Ala Ser Tyr Leu Phe Lys	
	100 105 110	

23

- Asn Arg Ile Val Tyr Leu Gly Met Ser Leu Val Pro Ser Val Thr Glu
115 120 125
- 5 Leu Ile Leu Ala Glu Phe Leu Tyr Leu Gln Tyr Glu Asp Glu Glu Lys
130 135 140
- 10 Pro Ile Tyr Leu Tyr Ile Asn Ser Thr Gly Thr Thr Lys Asn Gly Glu
145 150 155 160
- 15 Lys Leu Gly Tyr Asp Thr Glu Ala Phe Ala Ile Tyr Asp Val Met Gly
165 170 175
- 20 Tyr Val Lys Pro Pro Ile Phe Thr Leu Cys Val Gly Asn Ala Trp Gly
180 185 190
- Glu Ala Ala Leu Leu Leu Thr Ala Gly Ala Lys Gly Asn Arg Ser Ala
195 200 205
- 25 Leu Pro Ser Ser Thr Ile Met Ile Lys Gln Pro Ile Ala Arg Phe Gln
210 215 220
- 30 Gly Gln Ala Thr Asp Val Glu Ile Ala Arg Lys Glu Ile Lys His Ile
225 230 235 240
- 35 Lys Thr Glu Met Val Lys Leu Tyr Ser Lys His Ile Gly Lys Ser Pro
245 250 255
- Glu Gln Ile Glu Ala Asp Met Lys Arg Pro Lys Tyr Phe Ser Pro Thr
260 265 270
- 40 Glu Ala Val Glu Tyr Gly Ile Ile Asp Lys Val Val Tyr Asn Glu Arg
275 280 285
- 45 Gly Ser Gln Asp Arg Gly Val Val Ser Asp Leu Lys Lys Ala Gln Leu
290 295 300
- 50 Ile
305
- 55 <210> 17
<211> 906
<212> DNA
<213> Nicotiana tabacum
- 60 <220>
<221> CDS
<222> (45)..(755)
<223>
- <400> 17

24

	gcggccgctc caagattcat ccccaactct caacacattc aact atg cgc acc caa	56
	Met Arg Thr Gln	
	1	
5	att gtt cac aaa ctc ttt aac cga aga atc aac gga acc cct ttg aat	104
	Ile Val His Lys Leu Phe Asn Arg Arg Ile Asn Gly Thr Pro Leu Asn	
	5 10 15 20	
10	agt agt aag aga ctt tat ggg gta ata cca atg gta ata gag cac tct	152
	Ser Ser Lys Arg Phe Tyr Gly Val Ile Pro Met Val Ile Glu His Ser	
	25 30 35	
15	tca aga gga gaa agg gct tat gac ata ttc tca agg ota tta aag gaa	200
	Ser Arg Gly Glu Arg Ala Tyr Asp Ile Phe Ser Arg Leu Leu Lys Glu	
	40 45 50	
20	cga att att tgc att aac ggc ccc att gat gat tcc act tct cat gtt	248
	Arg Ile Ile Cys Ile Asn Gly Pro Ile Asp Asp Ser Thr Ser His Val	
	55 60 65	
25	gtt gtt gct cag ctt ctt ttt ctt gaa tct gag aac cct tct aag cct	296
	Val Val Ala Gln Leu Leu Phe Leu Glu Ser Glu Asn Pro Ser Lys Pro	
	70 75 80	
30	att cac aag tac ctc aac tct cca ggt ggc gct gtt aca gct ggt ctt	344
	Ile His Lys Tyr Leu Asn Ser Pro Gly Gly Ala Val Thr Ala Gly Leu	
	85 90 95 100	
35	gca atc tat gat acc acg cag tat atc cga tct cca act cat act ata	392
	Ala Ile Tyr Asp Thr Thr Gln Tyr Ile Arg Ser Pro Ile His Thr Ile	
	105 110 115	
40	tgc cta ggt caa gca gct tca atg gga tcc ctt ctc tta gct gca ggt	440
	Cys Leu Gly Gln Ala Ala Ser Met Gly Ser Leu Leu Leu Ala Ala Gly	
	120 125 130	
45	gca aag ggt gag aga cga tct ctc cct aat gct tca gtt atg att cac	488
	Ala Lys Gly Glu Arg Arg Ser Leu Pro Asn Ala Ser Val Met Ile His	
	135 140 145	
50	cag cct ttc ggt ggg tat agc ggg cag gct aaa gat ctg acg atc cac	536
	Gln Pro Phe Gly Gly Tyr Ser Gly Gln Ala Lys Asp Leu Thr Ile His	
	150 155 160	
55	aca aaa cag ata gtt cgg gta tgg gat act ttg aat gac cta tat gca	584
	Thr Lys Gln Ile Val Arg Val Trp Asp Thr Leu Asn Asp Leu Tyr Ala	
	165 170 175 180	
60	aag cat aca gga caa cct ata gaa ata att caa aag aat atg gat agg	632
	Lys His Thr Gly Gln Pro Ile Glu Ile Ile Gln Lys Asn Met Asp Arg	
	185 190 195	
65	gat tat ttc atg aca cct gaa gag gcg aag gag ttt gga ata atc gat	680
	Asp Tyr Phe Met Thr Pro Glu Glu Ala Lys Glu Phe Gly Ile Ile Asp	
	200 205 210	
70	gaa gtt ata gat gaa cga cca atg gct tta gta act gat gct gtt gca	728
	Glu Val Ile Asp Glu Arg Pro Met Ala Leu Val Thr Asp Ala Val Ala	
	215 220 225	
75	aat gaa gcc aaa gaa aaa ggt tca agc tagaaaaatt gctgtaatac	775
	Asn Glu Ala Lys Glu Lys Gly Ser Ser	
	230 235	

25

	tgatctcatt gcagcctttg ttagcattta ccacgcgtaa ctgctctcc attttactta	835
	ctggtgtatt tactthotag tattttatet gatgaggcga tacctcatta ctttgthtcc	895
5	tcagcggacg c	906
	<210> 18	
	<211> 237	
10	<212> PRT	
	<213> Nicotiana tabacum	
	<400> 18	
15	Met Arg Thr Gln Ile Val His Lys Leu Phe Asn Arg Arg Ile Asn Gly	
	1 5 10 15	
20	Thr Pro Leu Asn Ser Ser Lys Arg Phe Tyr Gly Val Ile Pro Met Val	
	20 25 30	
	Ile Glu His Ser Ser Arg Gly Glu Arg Ala Tyr Asp Ile Phe Ser Arg	
25	35 40 45	
	Leu Leu Lys Glu Arg Ile Ile Cys Ile Asn Gly Pro Ile Asp Asp Ser	
	50 55 60	
30	Thr Ser His Val Val Val Ala Gln Leu Leu Phe Leu Glu Ser Glu Asn	
	65 70 75 80	
35	Pro Ser Lys Pro Ile His Lys Tyr Leu Asn Ser Pro Gly Gly Ala Val	
	85 90 95	
40	Thr Ala Gly Leu Ala Ile Tyr Asp Thr Thr Gln Tyr Ile Arg Ser Pro	
	100 105 110	
	Ile His Thr Ile Cys Leu Gly Gln Ala Ala Ser Met Gly Ser Leu Leu	
45	115 120 125	
	Leu Ala Ala Gly Ala Lys Gly Glu Arg Arg Ser Leu Pro Asn Ala Ser	
	130 135 140	
50	Val Met Ile His Gln Pro Phe Gly Gly Tyr Ser Gly Gln Ala Lys Asp	
	145 150 155 160	
55	Leu Thr Ile His Thr Lys Gln Ile Val Arg Val Trp Asp Thr Leu Asn	
	165 170 175	
60	Asp Leu Tyr Ala Lys His Thr Gly Gln Pro Ile Glu Ile Ile Gln Lys	
	180 185 190	
	Asn Met Asp Arg Asp Tyr Phe Met Thr Pro Glu Glu Ala Lys Glu Phe	
	195 200 205	

	Gly	Ile	Ile	Asp	Glu	Val	Ile	Asp	Glu	Arg	Pro	Met	Ala	Leu	Val	Thr
	210						215					220				
5																
	Asp	Ala	Val	Ala	Asn	Glu	Ala	Lys	Glu	Lys	Gly	Ser	Ser			
	225					230					235					
10																
	<210>	19														
	<211>	447														
	<212>	DNA														
	<213>	Nicotiana tabacum														
15																
	<400>	19														
	gcggccgcgtt	goggacaaga	taatcgattc	aagagacaat	gtatttgaga	aaaggaacta									60	
	tgatgagata	ctcgcccaat	ctagagctat	gaggaaagcc	ggaccagggtg	ctcaggctgc									120	
20																
	tccatctggc	ttcaggtgac	tggaaagagcg	gtaatgggtcc	caaaccttca	ggaacaactg									180	
	ttgttccctt	atagtttcga	ggaacaaagt	tgtctggttac	ttgggtctgtg	ccggtataat									240	
25																
	gtaactggga	caaagaacat	attgtagaaa	ccttggttga	gctgtgaagt	ataggggttt									300	
	tacaactatt	atgcacaggt	ctgcaaagag	taccataat	gtcaattggg	tgtaccagta									360	
	tcaaacatc	agatagtgc	agtgtatgg	ataaatgaat	atagatctct	ctgatgtcat									420	
30																
	ttttctttta	tcatgttcag	cggcgcg												447	
	<210>	20														
35																
	<211>	996														
	<212>	DNA														
	<213>	Nicotiana tabacum														
	<400>	20														
40																
	gcggccgcgtt	gcggacaaga	taatcgattc	aagagacaat	gtatttgaga	aaaggaacta									60	
	tgatgagata	ctcgcccaat	ctagagctat	gaggaaagcc	ggaccagggtg	ctcaggctgc									120	
	tccatctggc	ttcaggtgac	tggaaagagcg	gtaatgggtcc	caaaccttca	ggaacaactg									180	
45																
	ttgttccctt	atagtttcga	ggaacaaagt	tgtctggttac	ttgggtctgtg	ccggtataat									240	
	gtaactggga	caaagaacat	attgtagaaa	ccttggttga	gctgtgaagt	ataggggttt									300	
	tacaactatt	atgcacaggt	ctgcaaagag	taccataat	gtcaattggg	tgtaccaggc									360	
50																
	ggcogctggc	ttcttctttg	cttctctctc	cgctttctag	ctcgacgggt	actgaaaatc									420	
	gogagctggg	ttctggtaaa	tcaactttca	tatccagttc	caatttctcc	tttgcaactt									480	
55																
	ctgttccacg	ttgcaggcca	aacggcggtc	gaggttattg	ttacagggtc	ccggtagcta									540	
	agtcttttga	ccatataccc	caaaaattca	gactggaaaa	tctcaaagat	ggactactgg									600	
60																
	acaactataa	aagtgccctt	cagtatcttt	acggccttag	tcttccacag	atggatatgt									660	
	tcatgacaga	agataaocca	gtacggcgac	agtcagaaa	cgccactgag	gatagtarat									720	
	ctgcggccgc	tggcagatgc	tccacgaarg	gataccacag	caggctcttg	ttagtccata									780	

27

	cacatogtat aatttatggc tgatagtggc cgtacgactt gcagtgttat tttgcaattt	840
5	cttttgttta atctacatat tgaactcttt cgtactactt attcaaaaac atgaaatcct	900
	gagcagacta gatgcatttg tttaatatca tgaatgcaag gaatccacct acagctgata	960
	tgtatacaaaa gatacctttt tttcaagagc ggcgcg	996
10	<210> 21	
	<211> 602	
	<212> DNA	
15	<213> Nicotiana tabacum	
	<220>	
	<221> CDS	
	<222> (2) .. (193)	
20	<223>	
	<400> 21	
25	g cgg cgg ctg gaa gat gtc cgg cgc caa gtc aac gaa gcg gct caa cct	49
	Arg Pro Leu Glu Asp Val Arg Arg Gln Val Asn Glu Ala Val Gln Pro	
	1 5 10 15	
	cgt cat aaa atc gac aag atg tat gtc gcc ttt act gac caa cca att	97
	Arg His Lys Ile Asp Lys Met Tyr Val Ala Phe Thr Asp Gln Pro Ile	
30	20 25 30	
	gag aag gtg caa cag tac act gaa agg gat cgt ttt ttg tct gtc tca	145
	Glu Lys Val Gln Gln Tyr Thr Glu Arg Asp Arg Phe Leu Ser Val Ser	
	35 40 45	
35	gag gcc atg gag ttt ggt ctc ata gat ggg gtg cta gaa aca gaa tac	193
	Glu Ala Met Glu Phe Gly Leu Ile Asp Gly Val Leu Glu Thr Glu Tyr	
	50 55 60	
40	tagttgcaaa tgaatottta gtagtacatg gtagctagcc ttccaatgac gaaaaagctg	253
	gtgttgctca ttaaccactt cgaagtacaa gaagctgggt cttgcaaatt tgtatcgtag	313
	aaatatctca actcttcaat ccaggaatgt ccaaaaagcct aattctgaag acgggttatag	373
45	aaagcgctct tgttttaacta tttttgtctc tectgcagat acactcagca cttttgtggg	433
	tattaatcag ggtcttaatt catcacttaa ccacaatcca gttggaagcg aagtgatcaa	493
	acacaaaagca gattcaggaa gatgtgtatt tttcccaaat atatattact ccaattgcta	553
50	tcatecccttc gctgtcgtta tgaaaggata tttatbttat agcggccgc	602
	<210> 22	
55	<211> 64	
	<212> PRT	
	<213> Nicotiana tabacum	
	<400> 22	
60	Arg Pro Leu Glu Asp Val Arg Arg Gln Val Asn Glu Ala Val Gln Pro	
	1 5 10 15	

28

Arg His Lys Ile Asp Lys Met Tyr Val Ala Phe Thr Asp Gln Pro Ile
20 25 30

5 Glu Lys Val Gln Gln Tyr Thr Glu Arg Asp Arg Phe Leu Ser Val Ser
35 40 45

10 Glu Ala Met Glu Phe Gly Leu Ile Asp Gly Val Leu Glu Thr Glu Tyr
50 55 60

15 <210> 23
<211> 16
<212> DNA
<213> artificial sequence

20 <220>
<223> primer
<400> 23
agaattcgcg gacgat

16

25 <210> 24
<211> 32
<212> DNA
<213> artificial sequence

30 <220>
<223> primer
<400> 24
ctcatgcggc cgcgcgcaac gcaattaatg tg

32

35 <210> 25
<211> 32
<212> DNA
<213> artificial sequence

40 <220>
<223> primer

45 <400> 25
tcattgcggc gcgagatcca gtccgatgta ac

32

50 <210> 26
<211> 21
<212> DNA
<213> artificial sequence

55 <220>
<223> primer

60 <400> 26
gtggattgat gtgatatccc c

21

<210> 27
<211> 21
<212> DNA
<213> artificial sequence

	<220>		
	<223>	primer	
5	<400>	27	
		gtaaggatct gagctacaca t	21
10	<210>	28	
	<211>	24	
	<212>	DNA	
	<213>	artificial sequence	
15	<220>		
	<223>	primer	
	<400>	28	
		tataaccatgg atttggcattc ttgtg	24
20	<210>	29	
	<211>	21	
	<212>	DNA	
	<213>	artificial sequence	
25	<220>		
	<223>	primer	
30	<400>	29	
		atagatctca cctggagcca g	21
35	<210>	30	
	<211>	19	
	<212>	DNA	
	<213>	artificial sequence	
40	<220>		
	<223>	primer	
	<400>	30	
		gagcccatgg caagaggag	19
45	<210>	31	
	<211>	22	
	<212>	DNA	
	<213>	artificial sequence	
50	<220>		
	<223>	primer	
	<400>	31	
		atagatctttt ctagcttgaa cc	22
55	<210>	32	
	<211>	21	
	<212>	DNA	
60	<213>	artificial sequence	
	<220>		
	<223>	primer	

BASF Aktiengesellschaft

20030949

PF 55117 EP

30

	<400> 32	
	tcagccatgg cccctggagg a	21
5	<210> 33	
	<211> 24	
	<212> DNA	
	<213> artificial sequence	
10	<220>	
	<223> primer	
	<400> 33	
	taagatcttc agtattctgt ttcc	24
15		

PCT/EP2004/013555

